

Problem 5.3.5

Random variables N and K have the joint PMF

$$P_{N,K}(n,k) = \begin{cases} \frac{(1-p)^{n-1}p}{n} & k=1,\dots,n; \\ 0 & \text{otherwise.} \end{cases} \rightarrow \text{k'ya g'one uniform}$$

Find the marginal PMFs $P_N(n)$ and $P_K(k)$.

$$P_N(n) = \sum_{k=1}^{\infty} P_{N,K}(n,k) = \sum_{k=1}^n \frac{(1-p)^{n-1}p}{n}$$

$$P_N(n) = (1-p)^{n-1}p$$

$n=1,2,\dots$

$N \sim \text{Geom}(p)$

$$P_K(k) = \sum_{n=1}^{\infty} \frac{(1-p)^{n-1}p}{n}$$

$$= p \sum_{n=1}^{\infty} \frac{A^{n-1}}{n} = \frac{p}{A} \sum_{n=1}^{\infty} \frac{A^n}{n}$$

$$p \frac{d}{dA} \left(\sum_{n=1}^{\infty} A^n \right) = p \sum_{n=1}^{\infty} n A^{n-1}$$

$$p \int \sum_{n=1}^{\infty} A^{n-1} dA = p \sum_{n=1}^{\infty} \frac{A^n}{n} = p A \sum_{n=1}^{\infty} \frac{A^{n-1}}{n}$$

$$= \int \frac{p}{1-A} dA = \frac{p}{(1-A)^2} = \frac{p}{p^2} = \frac{1}{p}, \quad k=1,2,\dots,n$$

uniform

Problem 5.4.2




Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 6e^{-(2x+3y)} & x \geq 0, y \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find $P[X > Y]$ and $P[X + Y \leq 1]$.

(b) Find $P[\min(X, Y) \geq 1]$.

(c) Find $P[\max(X, Y) \leq 1]$.




$$\int_{x=0}^{\infty} \int_{y=0}^x 6e^{-2x} e^{-3y} dy dx$$

$$= \int_{x=0}^{\infty} \left(-2e^{-3y} e^{-2x} \Big|_{y=0}^x \right) dx$$

$$-2e^{-5x} + 2e^{-2x}$$

$$= \int_{x=0}^{\infty} -2e^{-5x} + 2e^{-2x} dx$$

$$\left. \frac{2}{5}e^{-5x} - e^{-2x} \right|_{x=0}^{\infty} = -\frac{2}{5} + 1 = \frac{3}{5} //$$



$$\int_{x=0}^1 \int_{y=0}^x 6e^{-2x} e^{-3y} dy dx$$

$$\int_{x=0}^1 \left(-2e^{-2x} e^{-3y} \Big|_{y=0}^x \right) dx$$


$$\int_{x=0}^1 \left(-2e^{-5x} + 2e^{-2x} \right) dx$$

$$= \left. \frac{2}{5} e^{-5x} - e^{-2x} \right|_{x=0}^1 = \frac{2}{5} e^{-5} - e^{-2} - \frac{2}{5} + 1$$

$$= \frac{3}{5} + \frac{2}{5} e^{-5} - e^{-2}$$

b) $P[\min\{X, Y\} > 1]$


$$P[X > 1, Y > 1]$$



$$\int_{x=1}^{\infty} \int_{y=1}^{\infty} 6e^{-2x} e^{-3y} dy dx$$

$$\int_{x=1}^{\infty} \left(-2e^{-2x} e^{-3y} \Big|_{y=1}^{\infty} \right) dx$$


$$\int_{x=1}^{\infty} 2e^{-2x} e^{-3} dx = -e^{-2x} e^{-3} \Big|_{x=1}^{\infty} = e^{-5}$$

$$\begin{aligned}
 c) \quad & P[\max(X, Y) \leq 1] = P(X \leq 1, Y \leq 1) \\
 &= \int_{x=0}^1 \int_{y=0}^1 6e^{-2x} e^{-3y} dy dx \\
 &= \int_{x=0}^1 \left(2e^{-2x} e^{-3y} \Big|_{y=0}^1 \right) dx \\
 &= \int_{x=0}^1 -2e^{-2x} e^{-3} + 2e^{-2x} dx \\
 &= \left(e^{-2x} e^{-3} - e^{-2x} \right) \Big|_0^1 = e^{-5} - e^{-2} - e^{-3} + 1 \\
 &= 1 + e^{-5} - e^{-2} - e^{-3}
 \end{aligned}$$


Problem 5.5.5

X and Y are random variables with the joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 5x^2/2 & -1 \leq x \leq 1; \\ & 0 \leq y \leq x^2 \\ 0 & \text{otherwise.} \end{cases}$$




(a) What is the marginal PDF $f_X(x)$?

(b) What is the marginal PDF $f_Y(y)$?

$$a) \quad f_X(x) = \int_{y=-\infty}^{\infty} f_{X,Y}(x,y) dy$$

$$= \int_0^{x^2} \frac{5x^2}{2} dy = \frac{5x^2}{2} y \Big|_{y=0}^{x^2}$$

$$f_X(x) = \frac{5}{2} x^4, \quad -1 \leq x \leq 1$$




$$\begin{aligned}
 b) f_Y(y) &= \int_{-1}^{-\sqrt{y}} \frac{5x^2}{2} dx + \int_{+\sqrt{y}}^1 \frac{5x^2}{2} dx \\
 &= \left. \frac{5x^3}{6} \right|_{-1}^{-\sqrt{y}} + \left. \frac{5x^3}{6} \right|_{\sqrt{y}}^1 = \frac{5}{6} - \frac{5y\sqrt{y}}{6} \\
 &= -\frac{5y\sqrt{y}}{6} + \frac{5}{6}
 \end{aligned}$$

$$f_Y(y) = \frac{5}{3} - \frac{5}{3}y\sqrt{y} \quad 0 \leq y \leq 1$$

Example 5.8 Problem

Find the joint CDF $F_{X,Y}(x,y)$ when X and Y have joint PDF

$y \rightarrow v$
 $x \rightarrow u$



$$f_{X,Y}(x,y) = \begin{cases} 2 & 0 \leq y \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases} \quad (5.22)$$

(1) $x \leq 0$ veya $y \leq 0 \Rightarrow F_{X,Y}(x,y) = 0$

(2) $y \geq x$ ve $0 \leq x \leq 1$



$$\int_{v=0}^x \int_{u=v}^x 2 du dv = x^2$$



(3) $y < x$ ve $0 \leq x \leq 1$

$u = v \rightarrow x$

$v = 0 \rightarrow y$

$$\int_{v=0}^y \int_{u=v}^x 2 du dv = 2xy - y^2$$





$$F_{X,Y}(x,y) = \begin{cases} 0 & x < 0 \text{ veya } y < 0 \\ x^2 & x \leq y, 0 \leq x \leq 1 \\ 2xy - y^2 & y < x, 0 \leq x < 1 \\ 2y - y^2 & 1 \leq x, 0 \leq y \leq 1 \\ 1 & x > 1, y > 1 \end{cases}$$


Problem 5.5.9



X and Y have the joint PDF

$$f_{X,Y}(x,y) = \begin{cases} cy & 0 \leq y \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) Draw the region of nonzero probability.
- (b) What is the value of the constant c ?
- (c) What is $F_X(x)$?
- (d) What is $F_Y(y)$?
- (e) What is $P[Y \leq X/2]$?



b)

$$\iint_{S_{X,Y}} f_{X,Y}(x,y) dx dy = 1 \quad \text{olmalı}$$

$$\int_{y=0}^1 \int_{x=y}^1 cy dx dy = \int_{y=0}^1 \left(cyx \Big|_{x=y}^1 \right) dy$$

$$= \int_{y=0}^1 cy - cy^2 dy = \left. \frac{c}{2} y^2 - \frac{c}{3} y^3 \right|_0^1 = \frac{c}{2} - \frac{c}{3} = 1$$


$$\Rightarrow \boxed{c=6}$$

c) $F_X(x) = ?$

önce $f_X(x)$ 'i bulalım.

$$f_X(x) = \int f_{XY}(x,y) dy$$

$$f_X(x) = \int_{y=0}^x 6y dy, \quad 0 \leq x \leq 1$$



$$= 3y^2 \Big|_{y=0}^x = 3x^2 \Rightarrow \boxed{f_X(x) = 3x^2, \quad 0 \leq x \leq 1}$$

$$F_X(x) = \int_0^x f_X(u) du = \int_0^x 3u^2 du = u^3 \Big|_0^x = \boxed{x^3}$$

d) $F_Y(y) = \int_0^y f_Y(y) dy$

$$f_Y(y) = \int f_{XY}(x,y) dx \quad x = y \rightarrow 1$$


$$= \int_{x=y}^1 6y dx = 6y x \Big|_{x=y}^{x=1} = 6y - 6y^2 \quad 0 \leq y \leq 1$$

$$F_Y(y) = \int_0^y (6v - 6v^2) dv = 3v^2 - 2v^3 \Big|_{v=0}^y$$

$$\boxed{F_Y(y) = \begin{cases} 0 & y \leq 0 \\ 3y^2 - 2y^3 & 0 \leq y \leq 1 \\ 1 & 1 < y \end{cases}}$$

e) $P(Y < X/2)$

$X > 2Y$



$$P(Y < X/2) = \int_{y=0}^{0.5} \int_{x=2y}^1 6y \, dx \, dy$$

$$\int_{y=0}^{0.5} \left(6y \times \left(x=1 \right)_{x=2y} \right) dy = \int_{y=0}^{0.5} 6y - 12y^2 \, dy$$

$$3y^2 - 4y^3 \Big|_{y=0}^{0.5}$$


$$= \frac{3}{4} - \frac{4}{8} = \frac{2}{8} = \frac{1}{4} //$$

Problem 5.6.7

In terms of a positive constant k , random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} k + 3x^2 & -1/2 \leq x \leq 1/2, \\ & -1/2 \leq y \leq 1/2, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) What is k ?
- (b) What is the marginal PDF of X ?
- (c) What is the marginal PDF of Y ?
- (d) Are X and Y independent?



$$1 = \int_{y=-1/2}^{1/2} \int_{x=-1/2}^{1/2} (k + 3x^2) dx dy$$

$$\int_{y=-1/2}^{1/2} \left(kx + x^3 \bigg|_{x=-1/2}^{x=1/2} \right) dy$$

$$\int_{y=-1/2}^{1/2} \left(k + \frac{1}{4} \right) dy = ky + \frac{y}{4} \bigg|_{-1/2}^{1/2} = k + \frac{1}{4} = 1$$

$$k = 3/4$$


$$b) f_X(x) = \int_{y=-1/2}^{1/2} f_{X,Y}(x,y) dy$$

$$\int_{y=-1/2}^{1/2} \left(\frac{3}{4} + 3x^2 \right) dy = \left(\frac{3y}{4} + 3x^2 y \right) \bigg|_{y=-1/2}^{y=1/2}$$

$$f_X(x) = \frac{3}{4} + 3x^2, \quad -\frac{1}{2} \leq x \leq \frac{1}{2}$$

$$c) f_Y(y) = \int_{x=-1/2}^{1/2} f_{X,Y}(x,y) dx = \int_{x=-1/2}^{1/2} \left(\frac{3}{4} + 3x^2 \right) dx$$

$$= \left. \frac{3}{4}x + x^3 \right|_{-1/2}^{1/2} = \frac{3}{4} + \frac{1}{4} = 1, \quad -\frac{1}{2} \leq y \leq \frac{1}{2}$$



$$f_X(x) f_Y(y) = 1 \times \left(\frac{3}{4} + 3x^2 \right)$$

? $f_{XY}(x, y) = \frac{3}{4} + 3x^2$
 EVET, EŞİTİĞİ
 YANI BAĞIMSIZDIR

Problem 5.6.8

X_1 and X_2 are independent, identically distributed random variables with PDF

$$f_X(x) = \begin{cases} x/2 & 0 \leq x \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$

- Find the CDF, $F_X(x)$.
- What is $P[X_1 \leq 1, X_2 \leq 1]$, the probability that X_1 and X_2 are both less than or equal to 1?
- Let $W = \max(X_1, X_2)$. What is $F_W(1)$, the CDF of W evaluated at $w = 1$?
- Find the CDF $F_W(w)$.

a)
$$F_X(x) = \int_{u=0}^x f_X(u) du = \int_{u=0}^x \frac{u}{2} du = \left. \frac{u^2}{4} \right|_{u=0}^x$$

$$f_X(x) = \begin{cases} \frac{x^2}{4}, & 0 \leq x \leq 2 \\ 1 & 2 \leq x \\ 0 & x < 0 \end{cases}$$

b)
$$\begin{aligned} P[X_1 \leq 1, X_2 \leq 1] &= P[X_1 \leq 1] P[X_2 \leq 1] \\ &= F_{X_1}(1) F_{X_2}(1) = \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{16} \end{aligned}$$

$$c) W = \max(X_1, X_2)$$

$$F_W(1) = P(W \leq 1) = P(\max(X_1, X_2) \leq 1)$$

$$= P(X_1 \leq 1, X_2 \leq 1)$$

$$= P(X_1 \leq 1) P(X_2 \leq 1)$$

$$= F_{X_1}(1) F_{X_2}(1) = \frac{1}{16} \text{ aynı sayı}$$

$$d) F_W(w) = ?$$

$$= P(W \leq w)$$

$$= P(X_1 \leq w) P(X_2 \leq w)$$

$$= \frac{w}{2} \cdot \frac{w}{2} = \frac{w^2}{4}$$

$$F_W(w) = \begin{cases} 0 & w \leq 0 \\ \frac{w^2}{4} & 0 < w \leq 2 \\ 1 & 2 \leq w \end{cases}$$

Problem 5.7.5

X and Y are random variables with $E[X] = E[Y] = 0$ and $\text{Var}[X] = 1$, $\text{Var}[Y] = 4$ and correlation coefficient $\rho = 1/2$. Find $\text{Var}[X + Y]$.

$$\text{Covariance (Kovaryans)} = \text{Cov}(X, Y) \\ = E[XY] - E[X]E[Y]$$

$$\text{Correlation coefficient} = \rho = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

İnti Katsayısı

$$\rho = \frac{1}{2} = \frac{E[XY] - E[X]E[Y]}{\sqrt{\text{Var}(X)\text{Var}(Y)}} = \frac{E[XY]}{2}$$

$-1 \leq \rho \leq 1$

$$E[XY] = 1 \rightarrow \text{Cov}(X, Y) = 1$$

$$\text{Var}(X+Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$$

$$= E[(X+Y)^2] - (E[X+Y])^2$$

$$= E[X^2 + 2XY + Y^2] - (E[X] + E[Y])^2$$

$$= E[X^2] + 2E[XY] + E[Y^2] - (E[X]^2 + E[Y]^2 - 2E[X]E[Y])$$

$$= \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$$

$$\text{Var}(X+Y) = 1 + 4 + 2 = 7$$

$$E[(X - \mu_X)(Y - \mu_Y)] > 0$$



Si fueriati $\Rightarrow E[XY] = E[X]E[Y]$

$$\Downarrow$$

$$\text{Cov}(X, Y) = 0$$

Problem 5.7.9

X and Y are identically distributed random variables with $E[X] = E[Y] = 0$ and covariance $\text{Cov}[X, Y] = 3$ and correlation coefficient $\rho_{X, Y} = 1/2$. For nonzero constants a and b , $U = aX$ and $V = bY$.

- Find $\text{Cov}[U, V]$.
- Find the correlation coefficient $\rho_{U, V}$.
- Let $W = U + V$. For what values of a and b are X and W uncorrelated?

$$\begin{aligned}
 \text{a) } \text{Cov}(U, V) &= E[UV] - E[U]E[V] \\
 &= E[abXY] - E[aX]E[bY] \\
 &= abE[XY] - abE[X]E[Y] \\
 &= ab(E[XY] - E[X]E[Y]) \\
 &= ab \text{Cov}(X, Y) \\
 &= 3ab //
 \end{aligned}$$

$$b) \rho_{u,v} = \frac{\text{Cov}(u,v)}{\sqrt{\text{Var}(u) \text{Var}(v)}} = \frac{3ab}{\sqrt{a^2 \text{Var}(x) b^2 \text{Var}(y)}} = \frac{3 \rightarrow \text{Cov}(x,y)}{\sqrt{\text{Var}(x) \text{Var}(y)}} = \rho_{x,y} = \frac{1}{2}$$

$$c) W = u + v$$

$$W = aX + bY$$

$$E[XW] = \underbrace{E[X]}_0 E[W] \Rightarrow \text{hangi } a \text{ ve } b \text{ için?}$$

Yani:

$$E[XW] = 0 \Rightarrow \text{hangi } a \text{ ve } b \text{ için}$$

$$E[X(aX + bY)] = a \underbrace{E[X^2]}_{\text{Var}(x)=6} + b \underbrace{E[XY]}_3$$

$$= 6a + 3b = 0 \text{ olursa}$$

$$\text{Var}(x) = ?$$

$$\boxed{2a + b = 0} \text{ olursa}$$

$$\frac{3}{\sqrt{\text{Var}(x) \text{Var}(y)}} = \frac{1}{2}$$

X ve W ilişkili
olun.

$$6^2 = \text{Var}(x) \text{Var}(y)$$

$$\text{Var}(x) = 6 \rightarrow E[X^2] = 6$$

Problem 5.7.10

True or False: For identically distributed random variables Y_1 and Y_2 with $E[Y_1] = E[Y_2] = 0$, $\text{Var}[Y_1 + Y_2] \geq \text{Var}[Y_1]$.

$$\rightarrow \text{Var}(Y_1 + Y_2) = \underbrace{\text{Var}(Y_1)}_{\sigma^2} + \underbrace{\text{Var}(Y_2)}_{\sigma^2} + 2 \text{Cov}(Y_1, Y_2) \stackrel{?}{\geq} \underbrace{\text{Var}(Y_1)}_{\sigma^2}$$

$$\times \text{Cov}(Y_1, Y_2) = E(Y_1 Y_2) - \underbrace{E(Y_1)}_0 \underbrace{E(Y_2)}_0$$

$$\rightarrow \text{Var}(Y_2) + 2 \text{Cov}(Y_1, Y_2) \stackrel{?}{\geq} 0$$

$$\rightarrow \text{Var}(Y_2) + 2 \text{Cov}(Y_1, Y_2) \sqrt{\frac{\text{Var}(Y_1)}{\sigma^2} \frac{\text{Var}(Y_2)}{\sigma^2}} \geq 0$$
$$\sqrt{\text{Var}(Y_1)} \sqrt{\text{Var}(Y_2)}$$

$$\sigma^2 + 2\rho \sigma^2 \geq 0$$

$$1 + 2\rho \geq 0 \Rightarrow \rho \geq -\frac{1}{2}$$

Demek ki $-1 \leq \rho < -\frac{1}{2}$ olursa o zaman bu eşitlik sağlanmaz.
her zaman.

Problem 5.7.11

X and Y are random variables with $E[X] = E[Y] = 0$ such that X has standard deviation $\sigma_X = 2$ while Y has standard deviation $\sigma_Y = 4$.

(a) For $V = X - Y$, what are the smallest and largest possible values of $\text{Var}[V]$?

(b) For $W = X - 2Y$, what are the smallest and largest possible values of $\text{Var}[W]$?

$$a) \quad \text{Var}(V) = \text{Var}(X - Y) = \underbrace{\text{Var}(X)}_4 + \underbrace{\text{Var}(Y)}_{16} - 2 \text{Cov}(X, Y)$$

$$-1 \leq \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y} = \rho_{X,Y} \leq 1$$

$\sigma_X = 2$
 $\sigma_Y = 4$

$$-8 \leq \text{Cov}(X, Y) \leq 8$$

$$\text{Var}(V) = 20 - 2 \text{Cov}(X, Y)$$

$$4 \leq \text{Var}(V) \leq 36$$

$$b) \quad W = X - 2Y$$

$$\text{Var}(W) = \text{Var}(X - 2Y)$$

$$= E[(X - 2Y)^2] - (E[X - 2Y])^2$$

$$= E[X^2 - 4XY + 4Y^2] - (E[X])^2 + 4E[X]E[Y] - 4E[Y]^2$$

$$= E[X^2] - E[X]^2 + 4(E[Y^2] - E[Y]^2)$$

$$= \underbrace{\text{Var}(X)}_4 + 4 \underbrace{\text{Var}(Y)}_{16} - 4 \text{Cov}(X, Y) = \text{Var}(W)$$

$$= 68 - 4 \operatorname{Cov}(X, Y) = \operatorname{Var}(W)$$

$$= 68 - 4 \rho_{xy} \cdot 8 = 68 - 32 \rho_{xy} = \operatorname{Var}(W)$$

$$\boxed{36 \leq 68 - 32 \rho_{xy} \leq 100} //$$


Problem 5.7.13

Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 5x^2/2 & -1 \leq x \leq 1; \\ & 0 \leq y \leq x^2, \\ 0 & \text{otherwise.} \end{cases}$$

Answer the following questions.

- (a) What are $E[X]$ and $\operatorname{Var}[X]$?
 (b) What are $E[Y]$ and $\operatorname{Var}[Y]$?
 (c) What is $\operatorname{Cov}[X, Y]$?
 (d) What is $E[X + Y]$?
 (e) What is $\operatorname{Var}[X + Y]$?



a) Once $f_X(x)$ is found:

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy = \int_0^{x^2} \frac{5x^2}{2} dy = \frac{5x^2}{2} y \Big|_{y=0}^{x^2} = \frac{5x^4}{2}, \quad -1 \leq x \leq 1$$

$$E[X] = \int_{-1}^1 x \frac{5x^4}{2} dx = \frac{5x^6}{12} \Big|_{-1}^1 = 0 //$$

$$\operatorname{Var}(X) = E[X^2] = \int_{-1}^1 x^2 \frac{5x^4}{2} dx = \frac{5x^7}{14} \Big|_{-1}^1 = \frac{5}{7} //$$

$$\begin{aligned}
 b) \quad f_Y(y) &= \int_{-\infty}^{\infty} f_{X,Y}(x) dx \\
 &= \int_{-\sqrt{y}}^{-1} f_{X,Y}(x) dx + \int_{\sqrt{y}}^1 f_{X,Y}(x) dx \\
 &= \int_{-1}^{-\sqrt{y}} \frac{5x^2}{2} dx + \int_{\sqrt{y}}^1 \frac{5x^2}{2} dx \\
 &= \left. \frac{5x^3}{6} \right|_{-1}^{-\sqrt{y}} + \left. \frac{5x^3}{6} \right|_{\sqrt{y}}^1 = \left(\frac{5}{6} - \frac{5y\sqrt{y}}{6} \right) 2 \\
 &\quad 0 \leq y \leq 1
 \end{aligned}$$

$$\begin{aligned}
 E[Y] &= \int_0^1 \left(\frac{5y}{3} - \frac{5}{3} y^{3/2} \right) dy \\
 &= \left. \frac{5y^2}{6} - \frac{5}{3} \frac{y^{7/2}}{7/2} \right|_0^1 = \frac{5}{6} - \frac{5}{3} \frac{2}{7} \\
 &= \frac{5}{6} - \frac{10}{21} = \frac{35-20}{42} = \frac{15}{42}
 \end{aligned}$$

$$\begin{aligned}
 E[Y^2] &= \int_0^1 \left(\frac{10y^2}{3} - \frac{10}{3} y^{7/2} \right) dy = \left. \frac{10y^3}{9} - \frac{10}{3} \frac{y^{9/2}}{9/2} \right|_0^1 \\
 &= \frac{10}{9} - \frac{10}{3} \times \frac{2}{9} = \frac{10}{9} - \frac{20}{27} \\
 &= \frac{10}{27}
 \end{aligned}$$

$$\begin{aligned}
 \text{Var}(Y) &= \frac{10}{27} - \left(\frac{5}{14} \right)^2 \\
 &= 0.24
 \end{aligned}$$

$$\text{Cov}(X, Y) = E(XY) - \underbrace{E(X)}_0 \underbrace{E(Y)}_{5/14}$$

$$\begin{aligned} E[XY] &= \int_{-1}^1 \int_0^{x^2} xy \frac{5x^2}{2} dy dx = \int_{-1}^1 \int_0^{x^2} \frac{5x^3 y}{2} dy dx \\ &= \int_{-1}^1 \left(\frac{5x^3 y^2}{4} \bigg|_{y=0}^{y=x^2} \right) dx = \int_{-1}^1 \frac{5}{4} x^7 dx \\ &= \frac{5}{4} \frac{x^8}{8} \bigg|_{-1}^1 = 0 \end{aligned}$$

$$d) E[X+Y] = E[X] + E[Y] = 0 + \frac{5}{14} = \frac{5}{14}$$

$$\begin{aligned} e) \text{Var}(X+Y) &= \underbrace{\text{Var}(X)}_{5/7} + \underbrace{\text{Var}(Y)}_{0.24} + 2 \underbrace{\text{Cov}(X, Y)}_0 \\ &= 0.96 \end{aligned}$$

Problem 5.7.14

Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 2 & 0 \leq y \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) What are $E[X]$ and $\text{Var}[X]$?
- (b) What are $E[Y]$ and $\text{Var}[Y]$?
- (c) What is $\text{Cov}[X, Y]$?
- (d) What is $E[X + Y]$?
- (e) What is $\text{Var}[X + Y]$?



Problem 5.7.15

A transmitter sends a signal X and a receiver makes the observation $Y = X + Z$, where Z is a receiver noise that is independent of X and $E[X] = E[Z] = 0$. Since the average power of the signal is $E[X^2]$ and the average power of the noise is $E[Z^2]$, a quality measure for the received signal is the signal-to-noise ratio

$$\Gamma = \frac{E[X^2]}{E[Z^2]} = \frac{\text{signal gücü}}{\text{gürültü gücü}}$$

How is Γ related to the correlation coefficient $\rho_{X,Y}$?

$$\rho_{X,Y} = \frac{\text{Cov}(X,Y)}{\sqrt{\text{Var}(X) \cdot \text{Var}(Y)}} = \frac{\text{Cov}(X,Y)}{\sqrt{E[X^2]} \sqrt{\text{Var}(X+Z)}}$$

$$\rho_{X,Y} = \frac{\text{Cov}(X, X+Z)}{\sqrt{\text{Var}(X)} \sqrt{\text{Var}(X)+\text{Var}(Z)}}$$

X ile Z bağımsız
↓
ilintisi.

$$\begin{aligned} \text{Cov}(X, X+Z) &= E[X^2 + XZ] - E[X]E[X+Z] \\ &= E[X^2] - \underbrace{E[XZ]}_0 - E[X](E[X] + \underbrace{E[Z]}_0) \end{aligned}$$

$$\text{Cov}(X, X+Z) = E[X^2] - \underbrace{E[X]^2}_0 = \text{Var}(X) = E[X^2]$$


$$\rho_{X,Y} = \frac{\text{Var}(X)}{\sqrt{\text{Var}(X)} \sqrt{\text{Var}(X)+\text{Var}(Z)}} = \frac{\sqrt{\text{Var}(X)}}{\sqrt{\text{Var}(X)+\text{Var}(Z)}}$$

$$= \frac{1}{\sqrt{1 + \frac{\text{Var}(Z)}{\text{Var}(X)}}} = \frac{1}{\sqrt{1 + \frac{E[Z^2]}{E[X^2]}}} = \boxed{\frac{1}{\sqrt{1 + \frac{1}{\Gamma}}}} = \rho_{X,Y}$$

Yani: $\Gamma \uparrow \Leftrightarrow \rho_{X,Y} \rightarrow 1$

Problem 5.8.6

The random variables X and Y have joint PMF



Find

- The expected values $E[X]$ and $E[Y]$,
- The variances $\text{Var}[X]$ and $\text{Var}[Y]$,
- The correlation, $r_{X,Y} = E[XY]$,
- The covariance, $\text{Cov}[X, Y]$,
- The correlation coefficient, $\rho_{X,Y}$.

$$P_X = \left[\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4} \right]$$

$$P_Y = \left[\frac{25}{48}, \frac{13}{48}, \frac{7}{48}, \frac{3}{48} \right]$$

$$S_X = \{1, 2, 3, 4\}$$

$$S_Y = \{1, 2, 3, 4\}$$

$$a) \quad E[X] = \frac{10}{4} = \frac{5}{2} \quad E[Y] = \frac{25 + 26 + 21 + 12}{48} = \frac{84}{48} = \frac{7}{4}$$

$$b) \quad E[X^2] = \frac{1 + 4 + 9 + 16}{4} = \frac{15}{2} \quad E[Y^2] = \frac{25 + 52 + 63 + 48}{48} = \frac{188}{48} = \frac{47}{12}$$

$$\text{Var}(X) = \frac{15}{2} - \frac{25}{4} = \frac{5}{4}$$

$$\text{Var}(Y) = \frac{47}{12} - \frac{49}{16} = \frac{4 \cdot 47 - 3 \cdot 49}{48} = \frac{41}{48}$$

$$c) \quad r = E[XY]$$

$$= 1 \times \frac{1}{4} + 2 \times \frac{1}{8} + 4 \times \frac{1}{8} + \frac{3}{12} + \frac{6}{12} + \frac{9}{12} + \frac{4}{16} + \frac{8}{16} + \frac{12}{16} + \frac{16}{16}$$

$$= \frac{1}{4} + \frac{1}{4} + \frac{1}{2} + \frac{18}{12} + \frac{40}{16}$$

$$= 1 + \frac{72}{48} + \frac{120}{48} = 5$$

$$d) \text{Cov}(X, Y) = \underbrace{E(XY)}_r - \underbrace{E(X)}_{5/2} \underbrace{E(Y)}_{7/4}$$

$$= 5 - \frac{35}{8} = \frac{5}{8} > 0$$

$$e) \rho_{XY} = \frac{\text{Cov}(XY)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} = \frac{5/8}{\sqrt{\frac{5}{4} \frac{41}{48}}} = \frac{5}{8} \sqrt{\frac{192}{205}} = 0.6$$

Problem 5.8.8

Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 1/2 & -1 \leq x \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find $r_{X,Y}$ and $E[e^{X+Y}]$.

$$E[XY]$$


$$E[XY] = \int_{y=-1}^1 \int_{x=-1}^y xy \frac{1}{2} dx dy$$

$$\int_{y=-1}^1 \left(\frac{x^2 y}{4} \Big|_{x=-1}^y \right) dy = \int_{y=-1}^1 \left(\frac{y^3}{4} - \frac{y}{4} \right) dy$$

$$= \frac{y^4}{16} - \frac{y^2}{8} \Big|_{y=-1}^1 = \left(\frac{1}{16} - \frac{1}{8} \right) - \left(\frac{1}{16} - \frac{1}{8} \right) = 0$$

$$E[e^{X+Y}] = \int_{y=-1}^1 \int_{x=-1}^y e^x e^y \frac{1}{2} dx dy$$

$$= \int_{y=-1}^1 \left(\frac{e^x e^y}{2} \Big|_{x=-1}^y \right) dy$$



$$= \int_{y=-1}^1 e^{\frac{2y}{2}} - e^{\frac{y}{2}} e^{-1} dy = \left. \frac{e^{2y}}{4} - \frac{e^{\frac{y}{2}} e^{-1}}{2} \right|_{y=-1}^1$$

$$= \frac{e^2}{4} - \frac{e^{-2}}{4} - \frac{e^0}{2} + \frac{e^{-2}}{2}$$

$$E[e^{X+Y}] = \frac{e^2}{4} - \frac{1}{2} + \frac{e^{-2}}{4}$$

Problem 5.9.4

X_1 and X_2 are identically distributed Gaussian $(0,1)$ random variables. Moreover, they are jointly Gaussian. Under what conditions are X_1 , X_2 and $X_1 + X_2$ identically distributed?

$$f_{X,Y}(x,y) = \frac{\exp \left[- \frac{\left(\frac{x-\mu_x}{\sigma_x} \right)^2 - \frac{2\rho_{xy}(x-\mu_x)(y-\mu_y)}{\sigma_x\sigma_y} + \left(\frac{y-\mu_y}{\sigma_y} \right)^2}{2(1-\rho_{xy}^2)} \right]}{2\pi\sigma_x\sigma_y\sqrt{1-\rho_{xy}^2}}$$

$\rho_{xy}=0$ (yani ilintisiz) olduğunda

$$f_{X,Y}(x,y) = \exp \left[- \frac{\left(\frac{x-\mu_x}{\sigma_x} \right)^2}{2} - \frac{\left(\frac{y-\mu_y}{\sigma_y} \right)^2}{2} \right]$$

$$= \frac{1}{\sqrt{2\pi}\sigma_x} \exp \left(- \frac{(x-\mu_x)^2}{2\sigma_x^2} \right) \frac{1}{\sqrt{2\pi}\sigma_y} \exp \left(- \frac{(y-\mu_y)^2}{2\sigma_y^2} \right)$$

$f_X(x)$

$f_Y(y)$

yani X, Y bağımsızdır.

$$\mu_{x_1}, \mu_{x_2} = 0, \sigma_{x_1} = \sigma_{x_2} = 1$$

$$f_{x_1, x_2}(x_1, x_2) = \frac{\exp\left[-\frac{x_1^2 - 2\rho_{x_1, x_2}x_1x_2 + x_2^2}{2(1-\rho_{x_1, x_2}^2)}\right]}{2\pi\sqrt{1-\rho_{x_1, x_2}^2}}$$

$$x_1 \sim \mathcal{N}(0, 1)$$

$$x_2 \sim \mathcal{N}(0, 1)$$

$$x_1 + x_2 \sim \mathcal{N}(0, 2 + 2\rho_{x_1, x_2})$$

$$\begin{aligned}\text{Var}(x_1 + x_2) &= \underbrace{\text{Var}(x_1)}_1 + \underbrace{\text{Var}(x_2)}_1 + 2\underbrace{\text{Cov}(x_1, x_2)}_{\rho_{x_1, x_2} \cdot \cancel{\sigma_{x_1}} \cdot \cancel{\sigma_{x_2}} \cdot 1 \cdot 1} \\ &= 2 + 2\rho_{x_1, x_2}\end{aligned}$$

Eğer $2 + 2\rho_{x_1, x_2} = 1$ olursa o zaman $x_1 + x_2$ ve x_1, x_2

aynı dağılımdır

$$\boxed{\text{Yani } \rho_{x_1, x_2} = -\frac{1}{2} \text{ olursa}}$$

Problem 5.8.10

Random variables N and K have the joint PMF

$$P_{N,K}(n,k) = \begin{cases} (1-p)^{n-1}p/n & k=1, \dots, n; \\ & n=1, 2, \dots, \\ 0 & \text{otherwise.} \end{cases}$$

Find the marginal PMF $P_N(n)$ and the expected values $E[N]$, $\text{Var}[N]$, $E[N^2]$, $E[K]$, $\text{Var}[K]$, $E[N+K]$, $r_{N,K}$, $\text{Cov}[N, K]$.

$$P_N(n) = \sum_{k=1}^{\infty} \frac{(1-p)^{n-1}}{n} p = p \sum_{k=1}^{\infty} \frac{A^{n-1}}{n}$$

$$\sum_{n=1}^{\infty} (1-p)^n = \frac{1}{1-(1-p)} = \frac{1}{1-A}$$

$$\downarrow \frac{d}{dA}$$

$$\sum_{n=1}^{\infty} n(A)^{n-1}$$

Problem 5.9.5

Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = ce^{-(2x^2-4xy+4y^2)}.$$

(a) What are $E[X]$ and $E[Y]$? $\rightarrow \mu_x \text{ ve } \mu_y?$

(b) Find the correlation coefficient $\rho_{X,Y}$.

(c) What are $\text{Var}[X]$ and $\text{Var}[Y]$?

(d) What is the constant c ?

(e) Are X and Y independent?

$$f_{X,Y}(x,y) = \exp \left[\frac{\left(\frac{x-\mu_x}{\sigma_x} \right)^2 - \frac{2\rho_{xy}(x-\mu_x)(y-\mu_y)}{\sigma_x\sigma_y} + \left(\frac{y-\mu_y}{\sigma_y} \right)^2}{2(1-\rho_{xy}^2)} \right]$$

$$\frac{1}{2\pi\sigma_x\sigma_y\sqrt{1-\rho_{xy}^2}}$$

$$\begin{aligned}
 & \frac{x^2}{\sigma_x^2} - 2x \frac{\mu_x}{\sigma_x} + \frac{\mu_x^2}{\sigma_x^2} - \frac{2\rho_{xy}xy}{\sigma_x\sigma_y} + \frac{2\rho_{xy}x\mu_y}{\sigma_x\sigma_y} + \frac{2\rho_{xy}\mu_y\mu_x}{\sigma_x\sigma_y} \\
 & - \frac{2\rho_{xy}\mu_x\mu_y}{\sigma_x\sigma_y} + \frac{y^2}{\sigma_y^2} - \frac{2y\mu_y}{\sigma_y} + \frac{\mu_y^2}{\sigma_y^2}
 \end{aligned}$$

a)

$$= 2(1-\rho_{xy}^2)(2x^2 - 4xy + 4y^2)$$

$$\frac{\mu_x^2}{\sigma_x^2} - \frac{2\rho_{xy}\mu_x\mu_y}{\sigma_x\sigma_y} + \frac{\mu_y^2}{\sigma_y^2} = 0$$

$$2\mu_x^2 - 4\mu_x\mu_y + 4\mu_y^2 = 0$$

$$\mu_x^2 - 2\mu_x\mu_y + 2\mu_y^2 = 0$$

$$(\mu_x - \mu_y)^2 + \mu_y^2 = 0$$

$$\Rightarrow \mu_x = 0, \mu_y = 0$$

$$-2\frac{\mu_x}{\sigma_x} + \frac{2\rho_{xy}\mu_y}{\sigma_x\sigma_y} = 0$$

$$\mu_x = \rho_{xy} \frac{\mu_y}{\sigma_y} \Rightarrow \mu_x = 2\sigma_x \mu_y$$

$$\frac{2\rho_{xy}\mu_x}{\sigma_x\sigma_y} = \frac{2\mu_y}{\sigma_y}$$

$$\mu_x = \frac{\mu_y \sigma_x}{\rho_{xy}} \Rightarrow \mu_x = \frac{\mu_y}{2\sigma_y}$$

$$-2\rho_{xy}xy = -4xy$$

$\Downarrow$

$$\rho_{xy} = 2\sigma_x\sigma_y$$

$$\rho_{xy} = 2 \cdot \frac{1}{\sqrt{2}} \cdot \frac{1}{2}$$

$$\rho_{xy} = \frac{1}{\sqrt{2}} *$$


d) $C \Rightarrow \frac{1}{2\pi\sigma_x\sigma_y\sqrt{1-\rho_{xy}^2}} = \frac{1}{2\pi\frac{1}{\sqrt{2}}\frac{1}{2}\sqrt{1-\frac{1}{2}}} = \frac{2}{\pi} //$

e) $p_{x,y} \neq 0 \Rightarrow$ Bağımsız değil

Problem 5.9.6


An archer shoots an arrow at a circular target of radius 50 cm. The arrow pierces the target at a random position (X, Y) , measured in centimeters from the center of the disk at position $(X, Y) = (0, 0)$. The bullseye is a solid black circle of radius 2 cm, at the center of the target. Calculate the probability $P[B]$ of the event that the arrow hits the bullseye under each of the following models:

- ✓ (a) X and Y are iid continuous uniform $(-50, 50)$ random variables.
 (b) The PDF $f_{X,Y}(x, y)$ is uniform over the 50 cm circular target.
 (c) X and Y are iid Gaussian ($\mu = 0, \sigma = 10$) random variables.



$$f_X(x) = \frac{1}{\sqrt{2\pi}10} e^{-x^2/200}$$

$$f_Y(y) = \frac{1}{\sqrt{2\pi}10} e^{-y^2/200}$$



a) $f_X(x) = \frac{1}{100}, -50 \leq x \leq 50$

$$f_Y(y) = \frac{1}{100}, -50 \leq y \leq 50$$

Hitleri vurma $\Rightarrow X^2 + Y^2 \leq 4$

$$P(B) = P(X^2 + Y^2 \leq 4)$$

$$\int_{y=-2}^2 \int_{x=-\sqrt{4-y^2}}^{\sqrt{4-y^2}} \frac{1}{100} \frac{1}{100} dx dy$$

$$= \int_{-2}^2 \frac{2\sqrt{4-y^2}}{10000} dy = \int_{-2}^2 \frac{4\sqrt{1-(\frac{y}{2})^2}}{10000} dy$$

$$\sin \theta = \frac{y}{2}$$

$$\cos \theta = \frac{dy}{2}$$

$$P(B) = \frac{4}{1000} \int_{-\pi/2}^{\pi/2} \cos \theta \times 2 \cos \theta \, d\theta$$


$$= \frac{4}{1000} \int_{-\pi/2}^{\pi/2} \cos(2\theta) + 1 \, d\theta$$

$$2 \cos^2 \theta = \cos(2\theta) + 1$$

$$= \frac{4}{1000} \left. \frac{\sin(2\theta)}{2} + \theta \right|_{-\pi/2}^{\pi/2} = \frac{4\pi}{1000} \checkmark$$

b) $P(B) = \frac{\text{Siyah hedef alanı}}{\text{Daire alanı}} = \frac{\pi 2^2}{\pi 50^2} = \frac{4}{2500}$

c) $\boxed{\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ dx dy &= r \, dr \, d\theta \end{aligned}} \quad \begin{aligned} x^2 + y^2 &= r^2 \end{aligned}$



$$= \frac{x^2 + y^2}{200} \, dx \, dy$$

$$\Rightarrow \int_0^{2\pi} \int_0^2 \frac{1}{2\pi 100} e^{-\frac{r^2}{200}} r \, dr \, d\theta$$

Problem 5.10.7

X_1, X_2, X_3 are iid exponential (λ) random variables. Find:

- (a) the PDF of $V = \min(X_1, X_2, X_3)$,
 (b) the PDF of $W = \max(X_1, X_2, X_3)$.

θ

$$2r \, dr = du$$

$$\int_{\theta=0}^{2\pi} \left(\int_{u=0}^4 \frac{1}{200\sqrt{u}} e^{-u/200} \frac{du}{2} \right) d\theta$$

$$\int_{\theta=0}^{2\pi} \frac{1}{2\pi} (1 - e^{-4/200}) d\theta = 1 - e^{-4/200}$$

Problem 5.10.9

Random variables $X_1, X_2, \dots, X_n$ are iid; each X_j has CDF $F_X(x)$ and PDF $f_X(x)$. Consider

$$L_n = \min(X_1, \dots, X_n)$$

$$U_n = \max(X_1, \dots, X_n).$$

In terms of $F_X(x)$ and/or $f_X(x)$:

- Find the CDF $F_{U_n}(u)$.
- Find the CDF $F_{L_n}(l)$.
- Find the joint CDF $F_{L_n, U_n}(l, u)$.


$$\begin{aligned} a) F_{U_n}(u) &= P(U_n \leq u) = P(\max(X_1, \dots, X_n) \leq u) \\ &\stackrel{\text{independent}}{=} P(X_1 \leq u, X_2 \leq u, \dots, X_n \leq u) \\ &= \prod_{i=1}^n P(X_i \leq u) = \prod_{i=1}^n F_X(u) \\ &\stackrel{\text{identical dist.}}{=} (F_X(u))^n \end{aligned}$$

$$\begin{aligned} b) F_{L_n}(l) &= P(L_n \leq l) \\ &= P(\min(X_1, \dots, X_n) \leq l) \\ &= 1 - P(\min(X_1, \dots, X_n) > l) \\ &= 1 - P(X_1 > l, X_2 > l, \dots, X_n > l) \\ &= 1 - \prod_{i=1}^n (1 - F_X(l)) \end{aligned}$$

$$= 1 - (1 - F_X(l))^n$$

$$c) F_{L_n U_n}(l, u) = P(L_n \leq l, U_n \leq u)$$

$$= P(\min(X_1, \dots, X_n) \leq l, \max(X_1, \dots, X_n) \leq u)$$



Kırmızı : $P(\max(X_1, \dots, X_n) \leq u) = (F_X(u))^n$

Yeşil : $P(\max(X_1, \dots, X_n) \leq u, \min(X_1, \dots, X_n) \geq l)$

$$= P(l \leq X_1 \leq u, l \leq X_2 \leq u, \dots, l \leq X_n \leq u)$$

$$= \underbrace{P(l \leq X_1 \leq u)}_{F_X(u) - F_X(l)} \underbrace{P(l \leq X_2 \leq u)}_{\vdots} \dots \underbrace{P(l \leq X_n \leq u)}_{\vdots}$$

$$= (F_X(u) - F_X(l))^n$$

$$F_{U_n L_n}(l, u) = (F_X(u))^n - (F_X(u) - F_X(l))^n$$

Problem 5.10.6

The joint PMF of X , Y , and Z , the number of 1-page, 2-page, and 3-page downloads, respectively, in each of four downloads is given by

$$P_{X,Y,Z}(x,y,z) = \binom{4}{x,y,z} \frac{1}{3^x} \frac{1}{2^y} \frac{1}{6^z}$$

multinomial

- (a) In a group of four downloads, what is the PMF of the number of 3-page documents?
 (b) In a group of four downloads, what is the expected number of 3-page documents?
 (c) Given that there are two 3-page documents in a group of four, what is the joint PMF of the number of 1-page documents and the number of 2-page documents?
 (d) Given that there are two 3-page documents in a group of four, what is the expected number of 1-page documents?
 (e) In a group of four downloads, what is the joint PMF of the number of 1-page documents and the number of 2-page documents?

$\left(\frac{1}{3}\right)$: 1 page $\left(\frac{1}{2}\right)$: 2 page $\frac{1}{6}$: 3 page

a) Binom (3 page vega dequl)

$$P = \frac{1}{6} \quad n = 4$$

$$P_Z(z) = \binom{4}{z} \left(\frac{1}{6}\right)^z \left(\frac{5}{6}\right)^{4-z}$$

b) $E[Y] = 4 \times \frac{1}{6} = \frac{2}{3}$

c) Kalon 2 download ve bunlar ya 2 ya da 1 page.

$$P\left[\underset{\substack{\downarrow \\ X}}{1 \text{ page}}\right] = \frac{\frac{1}{3}}{\frac{5}{6}} = \boxed{\frac{2}{5}}$$

$$P\left[\underset{\substack{\downarrow \\ Y}}{2 \text{ page}}\right] = \frac{\frac{1}{2}}{\frac{5}{6}} = \boxed{\frac{3}{5}}$$

$$P_{X,Y}(x,y) = \binom{2}{x,y} \left(\frac{2}{5}\right)^x \left(\frac{3}{5}\right)^y \quad x,y \in \{0,1,2\} \quad x+y \leq 2 \quad \text{Binom}$$

$$P[1 \text{ page} | \text{not 3 page}] = \frac{P[1 \text{ page, not 3 page}]}{P[\text{not 3 page}]}$$

$$= \frac{P(1 \text{ page})}{P(\text{not 3 page})} = \frac{\frac{1}{3}}{\frac{5}{6}} = \frac{2}{5}$$

d) $E[X | 2 \text{ 3-page downloads}]$

$$= 2 \times \frac{2}{5} = \frac{4}{5}$$

e) 4 download $P(1 \text{ page}) = \frac{1}{3}$ $P(2 \text{ page}) = \frac{1}{2}$ $P(3 \text{ page}) = \frac{1}{8}$

$$P_{X,Y}(x,y) = \binom{4}{x,y,4-x-y} \left(\frac{1}{3}\right)^x \left(\frac{1}{2}\right)^y \left(\frac{1}{6}\right)^{4-x-y}$$

başta verilen olasılık ağırlık fonksiyonu ile aynı

Not: $\binom{2}{x,y} = \frac{2!}{x!y!}$

Problem 5.9.9

Your course grade depends on two test scores: X_1 and X_2 . Your score X_i on test i is Gaussian ($\mu = 74, \sigma = 16$) random variable, independent of any other test score. $\sigma^2 = 256$

(a) With equal weighting, grades are determined by $Y = X_1/2 + X_2/2$. You earn an A if $Y \geq 90$. What is $P[A] = P[Y \geq 90]$?

(b) A student asks the professor to choose a weight factor w , $0 \leq w \leq 1$, such that

$$Y = wX_1 + (1-w)X_2.$$

Find $P[A]$ as a function of the weight w . What value or values of w maximize $P[A] = P[Y \geq 90]$? $\frac{1}{2} = w$

(c) A different student proposes that the better exam is the one that should count and that grades should be based on $M = \max(X_1, X_2)$. In a fit of generosity, the professor agrees! Now what is $P[A] = P[M > 90]$?

(d) How generous was the professor? In a class of 100 students, what is the expected increase in the number of A's awarded?

a) $Y \sim \mathcal{N}(74, 8\sqrt{2})$

$$E[Y] = \frac{E[X]}{2} + \frac{E[X]}{2} = 74$$

$$\text{Var}(Y) = \frac{\text{Var}(X)}{4} + \frac{\text{Var}(X)}{4} = \frac{256}{4} \times 2 = 128$$

$$\sigma_Y = 8\sqrt{2}$$

$$P(Y \geq 90) = P\left(\frac{Y-74}{8\sqrt{2}} \geq \frac{90-74}{8\sqrt{2}}\right) = Q\left(\frac{16}{8\sqrt{2}}\right) = Q(\sqrt{2})$$

$$Q(A) = \int_A^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$$


$N(0,1)$

b) $Y = wX_1 + (1-w)X_2 \sim N\left(74, \sqrt{256w^2 + (1-w)^2 256}\right)$

$$E[Y] = 74$$

$$\begin{aligned} \text{Var}(Y) &= 256w^2 + (1-w)^2 256 \\ &= 256(w^2 + 1 - 2w + w^2) \\ &= 256(2w^2 - 2w + 1) \end{aligned}$$

$$\begin{aligned} P(Y \geq 90) &= P\left(\frac{Y-74}{8\sqrt{2}\sqrt{2w^2-2w+1}} \geq \frac{90-74}{8\sqrt{2}\sqrt{2w^2-2w+1}}\right) \\ &= Q\left(\frac{\sqrt{2}}{\sqrt{2w^2-2w+1}}\right) \end{aligned}$$



$\Rightarrow$ bunu maksimize etmek istiyoruz
 $\downarrow$
 A' 'yi minimize etmeli

$$\min \sqrt{\frac{2}{2w^2-2w+1}}$$

$$\Rightarrow \max \{2w^2 - 2w + 1\}$$

$$\begin{aligned} 4w - 2 &= 0 \\ w &= 1/2 \end{aligned}$$

c) $M = \max(X_1, X_2)$


$$\begin{aligned} P(M \geq 90) &= P(\max(X_1, X_2) \geq 90) \\ &= 1 - P(\max(X_1, X_2) < 90) \\ &= 1 - P(X_1 < 90) P(X_2 < 90) \end{aligned}$$

$$\begin{aligned}
 &= 1 - \left(\Phi \left(\frac{90-74}{16} \right) \right)^2 \\
 &= 1 - \left(1 - Q(1) \right)^2 \\
 \downarrow \quad Q(\sqrt{2}) &\geq 1 - (1 - Q(1))^2 \\
 100 Q(\sqrt{2}) &\geq 100 \left[1 - (1 - Q(1))^2 \right]
 \end{aligned}$$

Problem 6.1.5

Let X and Y be discrete random variables with joint PMF

$$P_{X,Y}(x,y) = \begin{cases} 0.01 & x = 1, 2, \dots, 10, \\ & y = 1, 2, \dots, 10, \\ 0 & \text{otherwise.} \end{cases}$$



What is the PMF of $W = \min(X, Y)$?

Problem 6.1.6

For random variables X and Y in Problem 6.1.5, what is the PMF of $W = \max(X, Y)$?

$$P(W=w) = P(\min(X, Y) = w)$$

$$= \frac{11-w}{100} + \frac{11-w}{100} - \frac{1}{100}$$

$$P(W=w) = \frac{21-2w}{100}$$

$$P(W=10) = \frac{1}{100} = P(X=10, Y=10)$$

$$P(W=1) = \frac{19}{100}$$


6.1.6

$$W = \max(X, Y)$$

$$\begin{aligned} P(W=w) &= P(\max(X, Y) = w) \\ &= P(X=w, Y \leq w) + P(X \leq w, Y=w) \\ &\quad - P(X=Y=w) \end{aligned}$$

$$\frac{w}{100} + \frac{w}{100} - \frac{1}{100}$$

$$P(W=w) = \frac{2w-1}{100}$$




Problem 6.2.6

Let X denote the position of the pointer after a spin on a wheel of circumference 1. For that same spin, let Y denote the area within the arc defined by the stopping position of the pointer:

$$\begin{aligned} 2\pi r &= 1 \\ r &= \frac{1}{2\pi} \end{aligned}$$

$$\begin{aligned} A &= \pi r^2 = \pi \frac{1}{4\pi^2} \\ A &= \frac{1}{4\pi} \end{aligned}$$



$$0 \leq X \leq 1$$

distributed.

$$0 \leq Y \leq \frac{1}{4\pi}$$

- What is the relationship between X and Y ?
- What is $F_Y(y)$?
- What is $f_Y(y)$?
- What is $E[Y]$?

$$a) \quad X = 1 \times \frac{\theta}{2\pi}$$

$$Y = \frac{1}{4\pi} \times \frac{\theta}{2\pi}$$

$$Y 4\pi = X$$

$$b) \quad F_Y(y) = P(Y \leq y) = P\left(\frac{X}{4\pi} \leq y\right) = P(X \leq 4\pi y)$$

$$\begin{aligned} F_Y(y) &= 4\pi y \\ 0 &\leq y \leq \frac{1}{4\pi} \end{aligned}$$

$$c) f_Y(y) = \frac{dF_Y(y)}{dy} = 4\pi, \quad 0 \leq y \leq \frac{1}{4\pi}$$

dargesten.

$$d) E[Y] = \frac{1}{8\pi}$$

Problem 6.2.8

X is the uniform $(0,1)$ random variable. Find a function $g(x)$ such that the PDF of $Y = g(X)$ is

$$f_Y(y) = \begin{cases} 3y^2 & 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

$$F_Y(y) = \begin{cases} y^3 & 0 \leq y \leq 1 \\ 1 & 1 < y \end{cases}$$

$$P(Y \leq y) = y^3 \quad 0 \leq y \leq 1$$

$$P(g(X) \leq y) = y^3 \quad 0 \leq y \leq 1$$

$$P(X \leq g^{-1}(y)) = y^3 \Rightarrow$$


$g(x)$ monotone is

$$g^{-1}(y) = y^{1/3}$$

$$g(x) = x^{1/3}$$


$$\left[\begin{array}{l} F_X(x) \\ U \sim \text{rand}(0,1) \Rightarrow F_X^{-1}(U) \sim f_X(x) \end{array} \right]$$

random sample generation



Problem 6.2.11

For the uniform $(0,1)$ random variable U , find the CDF and PDF of $Y = a + (b-a)U$ with $a < b$. Show that Y is the uniform (a,b) random variable.



$$\begin{aligned} P(Y \leq y) &= P(a + (b-a)U \leq y) \\ &= P\left(U \leq \frac{y-a}{b-a}\right) = \frac{y-a}{b-a}, \quad 0 < \frac{y-a}{b-a} < 1 \end{aligned}$$

$$F_Y(y) = \begin{cases} \frac{y-a}{b-a}, & a \leq y \leq b \\ 0 & \text{o.w.} \end{cases}$$

$\frac{d}{dy}$ ↓

$$f_Y(y) = \begin{cases} \frac{1}{b-a}, & a \leq y \leq b \\ 0 & \text{o.w.} \end{cases}$$

$$\left[Y = a + (b-a)U \rightarrow \begin{array}{l} a, b \text{ arası } U \text{ için} \\ \text{dağılımı} \end{array} \right]$$

0,1 arası U için dağılımı

$$-\ln(\text{rand}) \sim \text{expo}(1)$$


$$-\frac{1}{\lambda} \log(\text{rand}) \sim \text{expo}(\lambda)$$

Problem 6.3.2

In a 50 km cycling time trial, a rider's exact time T , measured in minutes, is the continuous uniform $(50, 60)$ random variable. However, a rider's recorded time R in seconds is obtained by rounding up T to next whole second. That is, if T is 50 minutes, 27.001 seconds, then $R = 3028$ seconds. On the other hand, if T is exactly 50 minutes 27 seconds, then $R = 3027$. What is the PMF of R ?

$$T = U(50, 60)$$

$\downarrow$
dk



R
 $\downarrow$
sn

$$\left[50 \text{ dk } 27,001 \text{ sn} \Rightarrow 50 \text{ dk } , 28 \text{ sn} \right]$$

3028 sn

$$P(R=r) = P\left(\left\lfloor \frac{r}{60} \right\rfloor < T \leq \left\lceil \frac{r}{60} \right\rceil\right), \quad 3000 \leq r \leq 3600$$




$\begin{array}{r} 3281 \overline{) 60} \\ 300 \\ \underline{281} \\ 240 \\ \underline{41} \end{array} \Rightarrow P(54 < T \leq 55) = \frac{1}{600}$

Problem 6.3.7

The input voltage to a rectifier is the continuous uniform $(0, 1)$ random variable U . The rectifier output is a random variable W defined by

$$W = g(U) = \begin{cases} 0 & U < 0, \\ U & U \geq 0. \end{cases}$$

Find the CDF $F_W(w)$ and the expected value $E[W]$.




$$P(W=0) = P(U < 0) = \frac{1}{2}$$

$$P(W \leq w) = P(U \leq w)$$


$$P(W \leq w) = 0, \quad w < 0$$

$$P(W=0) = \frac{1}{2}$$



$$F_W(w) = \begin{cases} 0 & w < 0 \\ \frac{1}{2} + \frac{w}{2} & 0 \leq w < 1 \\ 1 & 1 \leq w \end{cases}$$

$$f_w(w) = \frac{d}{dw} F_w(w) = \begin{cases} \frac{1}{2} \delta(w) + \frac{1}{2} & , 0 \leq w \leq 1 \\ 0 & , \text{o.w.} \end{cases}$$



$$\begin{aligned} E[w] &= \int_0^1 w f_w(w) dw = \int_0^1 w \left(\frac{1}{2} \delta(w) + \frac{1}{2} \right) dw \\ &= \underbrace{\int_0^1 \frac{w}{2} \delta(w) dw}_0 + \underbrace{\frac{w^2}{4} \Big|_0^1}_{\frac{1}{4}} \end{aligned}$$

$$E[w] = \frac{1}{4}$$

Problem 6.3.10


The current X across a resistor is the continuous uniform $(-2, 2)$ random variable. The power dissipated in the resistor is $Y = 9X^2$ Watts.

(a) Find the CDF and PDF of Y .

(b) A power measurement circuit is range-limited so that its output is


$$W = \begin{cases} Y & Y < 16, \\ 16 & \text{otherwise.} \end{cases}$$

Find the PDF of W .





$$\begin{aligned} P(Y \leq y) &= P(9X^2 \leq y) = P\left(-\frac{\sqrt{y}}{3} \leq X \leq \frac{\sqrt{y}}{3}\right) \\ &= \int_{-\frac{\sqrt{y}}{3}}^{\frac{\sqrt{y}}{3}} \frac{1}{4} dy = \frac{2\sqrt{y}}{12} = \frac{\sqrt{y}}{6}, \quad 0 \leq y \leq 36 \end{aligned}$$

$$F_Y(y) = \begin{cases} 0 & y < 0 \\ \frac{\sqrt{y}}{6} & 0 \leq y \leq 36 \\ 1 & 36 < y \end{cases}$$



$$f_Y(y) = \frac{d}{dy} F_Y(y) = \begin{cases} \frac{1}{12\sqrt{y}} & , 0 \leq y \leq 36 \\ 0 & \text{o.w.} \end{cases}$$




$$S_W = [0, 16] \text{ mixed.}$$

$$P(W \leq w) = P(Y \leq w) \quad , \quad w < 16$$

$$= \frac{\sqrt{w}}{6} \quad , \quad 0 \leq w < 16$$

$$P(W \leq w) = 1 \quad 16 \leq w.$$



$$F_W(w) = \begin{cases} 0 & w \leq 0 \\ \sqrt{w}/6 & 0 < w < 16 \\ 1 & w \leq 16 \end{cases}$$

$$f_w(w) = \frac{d}{dw} F_w(w) = \begin{cases} \frac{1}{12\sqrt{w}} + \frac{1}{3} \delta(w-16) & , w > 0 \\ 0 & , w \leq 0 \end{cases}$$

$$E[W] = \int_0^{16} \frac{w}{12\sqrt{w}} + \frac{w}{3} \delta(w-16) dw$$

$$= \int_0^{16} \frac{\sqrt{w}}{12} dw + \int_0^{16} \frac{w}{3} \delta(w-16) dw$$

$$\frac{1}{18} w^{3/2} \Big|_0^{16} + \underline{\frac{16}{3}} = \frac{64}{18} + \frac{16}{3} = \frac{32}{9} + \frac{16}{3} = \frac{80}{9}$$

HATIRLA

$$\left[\begin{aligned} \int_{-\infty}^{\infty} g(x) \delta(x-x_0) dx &= \int_{-\infty}^{\infty} g(x_0) \delta(x-x_0) dx \\ &= g(x_0) \underbrace{\int_{-\infty}^{\infty} \delta(x-x_0) dx}_1 \\ &= g(x_0) \end{aligned} \right]$$

Problem 6.4.5

Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 6y & 0 \leq y \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let $W = Y - X$.

(a) What is S_W , the range of W ?

(b) Find $F_W(w)$ and $f_W(w)$.

$$W = Y - X$$

a) S_W range, deger analig

$$W = Y - X$$

$$Y \leq X \Rightarrow Y - X \leq 0$$

$$X=1 \text{ olabli } Y=0 \text{ olabli } \Rightarrow Y - X = -1$$

$$S_X = [-1, 0] \Rightarrow -1 \leq W \leq 0$$

$$b) P(W \leq w) = P(Y - X \leq w) = P(Y \leq X + w)$$

$$Y = X + w$$

$$w > 0 \text{ i\u00e7in } P(Y - X \leq w) = 1$$

$$-1 < w < 0 \text{ i\u00e7in}$$

$$P(Y - X \leq w)$$

$$= \int_{x=-w}^1 \int_{y=0}^{x+w} 6y \, dy \, dx$$



$$\int_{x=-w}^1 \left(3y^2 \Big|_0^{x+w} \right) dx = \int_{x=-w}^1 (3x^2 + 6xw + 3w^2) dx$$

$$= \left[x^3 + 3x^2w + 3w^2x \right]_{x=-w}^1$$

$$= (1 + 3w + 3w^2) - (-w^3 + 3w^3 - 3w^3)$$

$$P(W \leq w) = P(1 - X \leq w) = w^3 + 3w^2 + 3w + 1, \quad -1 \leq w < 0$$

$$w < -1 \Rightarrow P(W \leq w) = 0$$


$$F_W(w) = \begin{cases} 1 & w > 0 \\ w^3 + 3w^2 + 3w + 1 & -1 \leq w \leq 0 \\ 0 & w \leq -1 \end{cases}$$

$(w+1)^3$


$$(w+1)^3 = (w^2 + 2w + 1)(w+1)$$

$$= w^3 + 2w^2 + w + w^2 + 2w + 1$$

$$= w^3 + 3w^2 + 3w + 1$$




$$f_W(w) = \frac{d}{dw} F_W(w) = 3(w+1)^2, \quad -1 \leq w \leq 0$$



Problem 6.4.8

In a simple model of a cellular telephone system, a portable telephone is equally likely to be found anywhere in a circular cell of radius 4 km. (See Problem 5.5.4.) Find the CDF $F_R(r)$ and PDF $f_R(r)$ of R , the distance (in km) between the telephone and the base station at the center of the cell.



$$F_R(r) = P(R < r)$$

$$= \frac{\pi r^2}{\pi D^2}$$

$$F_R(r) = r^2 / D^2$$

$$f_R(r) = \frac{2r}{D^2}$$

$$\left[\frac{r^2}{D^2} = U \Rightarrow r = \sqrt{U D^2} = D \sqrt{U} = D \sqrt{\text{rand}} \right]$$

uniform(0,1) "rand"

$$\theta = 2\pi * \text{rand}$$




Problem 6.4.9

X and Y are independent identically distributed Gaussian $(0, 1)$ random variables. Find the CDF of $W = X^2 + Y^2$.

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \quad f_Y(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2} \quad -\infty < x, y < \infty$$

$$F_W(w) = P(W \leq w) = \iint_{X^2 + Y^2 \leq w} \frac{1}{2\pi} e^{-(x^2+y^2)/2} dy dx$$

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ dx dy &= r dr d\theta \end{aligned} \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \begin{array}{l} \text{Kutupsal koordinat} \\ \text{sistemiine geqis} \end{array}$$



$$\begin{aligned} \int_{\theta=0}^{2\pi} \int_{r=-\sqrt{w}}^{\sqrt{w}} \frac{1}{2\pi} e^{-r^2/2} r dr d\theta &= \int_{r=-\sqrt{w}}^{\sqrt{w}} \left(\frac{1}{2\pi} e^{-r^2/2} r \theta \right) \Big|_0^{2\pi} dr \\ &= \int_{r=-\sqrt{w}}^{\sqrt{w}} e^{-r^2/2} r dr \end{aligned}$$

$$\begin{aligned} r^2 &= u \quad 2r dr = du \Rightarrow \\ P(X^2 + Y^2 \leq w) &= \int_{u=0}^w e^{-u/2} \frac{1}{2} du = \left. -e^{-u/2} \right|_0^w \\ F_W(w) &= \boxed{P(W \leq w) = 1 - e^{-w/2}} \end{aligned}$$

$$f_W(w) = \frac{1}{2} e^{-w/2}, \quad w > 0 \Rightarrow \text{exponential} \left(\frac{1}{2} \right) \text{ istel}$$

$$\left(\text{Not: } \left(\frac{X}{\sqrt{2}} \right)^2 + \left(\frac{Y}{\sqrt{2}} \right)^2 = \frac{X^2 + Y^2}{2} \sim \text{istel}(1) \right)$$

Problem 6.4.11

X is the Gaussian $(0, 1)$ random variable and Z , independent of X , has PMF

$$P_Z(z) = \begin{cases} 1-p & z = -1, \\ p & z = 1. \end{cases}$$


$$Z = -1 \Rightarrow Y = -X \sim \mathcal{N}(0, 1)$$

$$Z = 1 \Rightarrow Y = X \sim \mathcal{N}(0, 1)$$


Find the PDF of $Y = ZX$.

$$f_Y(y)$$

$$f_Y(y) = f_{Y|Z}(y|z=-1) P(Z=-1) + f_{Y|Z}(y|z=+1) P(Z=+1)$$



$$X \rightarrow -1 \Rightarrow$$




$$f_Y(y) = f_X(x)$$

CDF:

$$P(Y \leq y) = P(X \leq y, Z=+1) + P(X \geq -y, Z=-1)$$

$$= P(X \leq y) P(Z=1) + P(X \geq -y) P(Z=-1)$$

$$= p \int_{-\infty}^y \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx + (1-p) \int_{-y}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$$




$$F_Y(y) = p F_X(y) + (1-p) F_X(y)$$

$$F_Y(y) = F_X(y) \quad \text{Y ve X aynı dağılımlı}$$

ortaman

$$f_Y(y) = f_X(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2}$$



$$N(\mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

Problem 6.4.13


For a constant $a > 0$, random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 1/a^2 & 0 \leq x, y \leq a, \\ 0 & \text{otherwise.} \end{cases}$$

Find the CDF and PDF of random variable

$$\rightarrow W = \max\left(\frac{X}{Y}, \frac{Y}{X}\right).$$

Hint: Is it possible to observe $W < 1$? (HAYIR)




$$\infty > W \geq 1$$

$$P(M \leq m) = P\left(\frac{X}{Y} \leq m\right)$$


$$P(X \leq mY)$$

$$m > 1: \frac{1}{a^2} \left(\left(a + a - \frac{a}{m} \right) \frac{a}{2} \right) = \frac{a^2}{a^2} \left(2 - \frac{1}{m} \right) \frac{1}{2} = 1 - \frac{1}{2m}$$




$$m < 1: \frac{a \times am}{2} \frac{1}{a^2} = \frac{m}{2}$$

$$F_M(m) = \begin{cases} \frac{m}{2} & 0 \leq m < 1 \\ 1 - \frac{1}{2m} & 1 \leq m \end{cases}$$




$$P(N \leq n) = P\left(\frac{Y}{X} \leq n\right) = P(Y \leq nX)$$

$$= \frac{a \cdot an}{2a^2} = \frac{n}{2}$$



$$n > 1: P(N \leq n) = \frac{(a + a - \frac{a}{n})a}{2a^2} \\ = \left(2 - \frac{1}{n}\right) \frac{1}{2} = 1 - \frac{1}{2n}$$



$$P(N < n) = \begin{cases} \frac{n}{2} & 0 \leq n < 1 \\ 1 - \frac{1}{2n} & n > 1 \end{cases}$$

$$W = \max\left(M, \frac{1}{M}\right)$$

$$P(W \leq w) = P\left(M \leq w, \frac{1}{M} \leq w\right)$$

W > 1

$$= P\left(M \leq w, \frac{1}{M} \leq w\right)$$

$$F_w(w) = 1 - \frac{1}{2w}, \quad w > 1$$

$$f_w(w) = \frac{1}{2w^2}, \quad w > 1$$

Problem 6.5.3

Find the PDF of $W = X + Y$ when X and Y have the joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 2 & 0 \leq x \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

$$P(W \leq w) = P(X + Y \leq w)$$

$$0 \leq W \leq 2 \Rightarrow S_W = [0, 2]$$

• $0 \leq w \leq 1$: Tarah alana göre:

$$P(W \leq w) = \underbrace{\left(\frac{w}{\sqrt{2}}\right)^2}_{\text{üçgen alan}} \underbrace{\frac{1}{2}}_{\text{pdf.}} \times 2 = \frac{w^2}{2}$$



• $1 \leq w \leq 2$:

$$P(W \leq w) = \left(\frac{1}{2} - \left(\frac{2-w}{\sqrt{2}}\right)^2 \frac{1}{2} \right) 2$$

$$1 - \frac{(2-w)^2}{2} = 1 - \frac{4 - 4w + w^2}{2}$$


$$= \frac{2 - 4 + 4w - w^2}{2}$$

$$P(W \leq w) = \frac{4w - w^2 - 2}{2} \quad 1 \leq w \leq 2$$



$$F_W(w) = \begin{cases} 0 & w < 0 \\ w^2/2 & 0 \leq w \leq 1 \\ \frac{4w - w^2 - 2}{2} & 1 \leq w \leq 2 \\ 1 & 2 < w \end{cases}$$

$$f_W(w) = \begin{cases} w & 0 \leq w \leq 1 \\ 2-w & 1 \leq w \leq 2 \\ 0 & \text{o.w.} \end{cases}$$




Problem 6.5.5

Random variables X and Y are independent exponential random variables with expected values $E[X] = 1/\lambda$ and $E[Y] = 1/\mu$. If $\mu \neq \lambda$, what is the PDF of $W = X + Y$? If $\mu = \lambda$, what is $f_W(w)$?

$$X \geq 0$$

$$Y \geq 0$$

$$F_W(w) = P(W \leq w) \\ = P(X+Y \leq w)$$



$$\int_{x=0}^w \int_{y=0}^{w-x} f_X(x) f_Y(y) dy dx$$

$$= \int_{x=0}^w \int_{y=0}^{w-x} \lambda e^{-\lambda x} \mu e^{-\mu y} dy dx$$

$$= \int_{x=0}^w \lambda e^{-\lambda x} \underbrace{\int_{y=0}^{w-x} \mu e^{-\mu y} dy}_{\text{red bracket}} dx$$

$$\int_{x=0}^w \lambda e^{-\lambda x} \left(1 - e^{-\mu(w-x)}\right) dx$$

$$= \int_{x=0}^w \lambda e^{-\lambda x} - \lambda e^{-\mu w} e^{-\lambda x} e^{\mu x} dx = \int_{x=0}^w \lambda e^{-\lambda x} - \lambda e^{-\mu w} e^{-(\lambda-\mu)x} dx$$

$$= \left[-e^{-\lambda x} + \frac{e^{-\mu w}}{\lambda - \mu} e^{-(\lambda-\mu)x} \right]_0^w$$

$$= -e^{-\lambda w} + \frac{e^{-\mu w}}{\lambda - \mu} e^{-(\lambda-\mu)w} + 1 - \frac{e^{-\mu w}}{\lambda - \mu}$$

$$F_W(w) = (1 - e^{-\lambda w}) - \frac{e^{-\mu w}}{\lambda - \mu} \left(1 - e^{-(\lambda-\mu)w}\right)$$

$\mu = \lambda$ also holds:

$$F_W(w) = 1 - e^{-\lambda w} \Rightarrow \text{exponential}$$

$$= -e^{-\lambda w} + \frac{\lambda}{\lambda-\mu} e^{-\lambda w} + 1 - \frac{\lambda}{\lambda-\mu} e^{-\mu w}$$

$$= \frac{\mu}{\lambda-\mu} e^{-\lambda w} - \frac{\mu}{\lambda-\mu} e^{-\mu w}$$

$$\int_0^w f_X(x) f_Y(w-x) dx = f_W(w)$$

$$= \int_0^w \lambda e^{-\lambda x} \mu e^{-\mu(w-x)} dx$$

$$= \int_0^w \lambda \mu e^{-\lambda x} e^{-\mu w} e^{\mu x} dx = \int_0^w \lambda \mu e^{-\mu w} e^{-(\lambda+\mu)x} dx$$

$$= \left(\frac{-\lambda \mu e^{-\mu w}}{\lambda+\mu} e^{-(\lambda+\mu)x} \right) \Big|_0^w$$

$$= \left(1 - e^{-(\lambda+\mu)w} \right) \frac{\lambda \mu}{\lambda+\mu} e^{-\mu w}$$

$$= \left(e^{-\mu w} - e^{-\lambda w} \right) \left(\frac{\lambda \mu}{\lambda+\mu} \right)$$

$\lambda = \mu$ i.d.R.

$$\int_0^w \lambda^2 e^{-\lambda x} e^{-\lambda(w-x)} dx = \int_0^w \lambda^2 e^{-\lambda w} dx$$

$$= \lambda^2 w e^{-\lambda w}$$

Erlang(2, λ)

2 i. d. r.d. in Toplam, Erlang 2 Lagrange.

5.3.6

$$P_{N,k}(n,k) = \begin{cases} \frac{100^n e^{-100}}{(n+1)!} & k=0,1,\dots,n \\ 0 & n=0,1,\dots \\ & 0 \leq k \leq n \end{cases}$$

$$P_N(n) = ?$$

$$P_N(n) = \sum_{k=0}^n P_{N,k}(n,k)$$

$$P_N(n) = \frac{100^n e^{-100}}{n!} \quad \text{Poisson}(100)$$

$$\begin{aligned} P_k(k) &= \sum_{n=0}^{\infty} P_{N,k}(n,k) = \sum_{n=0}^{\infty} \frac{100^n e^{-100}}{(n+1)!} \\ &= \sum_{n=0}^{\infty} \frac{1}{100} \frac{100^{n+1} e^{-100}}{(n+1)!} \end{aligned}$$

$$P_k(k) = \frac{1}{100} \left[1 - e^{-100} \right]$$

Problem 7.1.5

Every day you consider going jogging. Before each mile, including the first, you will quit with probability q , independent of the number of miles you have already run. However, you are sufficiently decisive that you never run a fraction of a mile. Also, we say you have run a marathon whenever you run at least 26 miles.

- ✓ (a) Let M equal the number of miles that you run on an arbitrary day. Find the PMF $P_M(m)$.
- ✓ (b) Let r be the probability that you run a marathon on an arbitrary day. Find r .
- ✓ (c) Let J be the number of days in one year (not a leap year) in which you run a marathon. Find the PMF $P_J(j)$. This answer may be expressed in terms of r found in part (b).
- (d) Define $K = M - 26$. Let A be the event that you have run a marathon. Find $P_{K|A}(k)$.

$$a) P_M(m) = (1-q)^m q, \quad m = 0, 1, \dots$$

$$\begin{aligned}
 b) P(\text{marathon}) &= P(M \geq 26) \\
 &= \sum_{m=26}^{\infty} (1-q)^m q \\
 &= (1-q)^{26} q \sum_{i=0}^{\infty} (1-q)^i \\
 &= (1-q)^{26} q \frac{1}{1-(1-q)} = (1-q)^{26} = r
 \end{aligned}$$

$$c) P_J(j) = \binom{365}{j} r^j (1-r)^{365-j}, \quad j = 0, 1, \dots, 365$$

Binom

$$d) P_{K|A}(k) = \begin{cases} \frac{P_K(k)}{P(A)}, & \text{if } k \in A \\ 0 & \text{o.w.} \end{cases}$$

→ A: Maraton koştuk. $\Rightarrow M \geq 26 \Rightarrow r$ olasılık

→ $K = M - 26 \rightarrow P_K(k) = P_M(k+26), k=0,1,\dots$ $\rightarrow P(A)=r$

$$P_{K|A}(k) = \begin{cases} \frac{P_K(k)}{r} & k=0,1,\dots,\infty \\ 0 & \text{o.w.} \end{cases}$$

$$= P_{K|A}(k) = \begin{cases} \frac{P_M(k+26)}{r} & k=0,1,\dots,\infty \\ 0 & \text{o.w.} \end{cases}$$

$$\frac{(1-q)^{k+26} q}{(1-q)^{26}} = (1-q)^k q$$

$$= P_{K|A}(k) = \begin{cases} (1-q)^k q & k=0,1,\dots,\infty \\ 0 & \text{o.w.} \end{cases}$$

Problem 7.1.7

A test for diabetes is a measurement X of a person's blood sugar level following an overnight fast. For a healthy person, a blood sugar level X in the range of 70 – 110 mg/dl is considered normal. When a measurement X is used as a test for diabetes, the result is called positive (event T^+) if $X \geq 140$; the test is negative (event T^-) if $X \leq 110$, and the test is ambiguous (event T^0) if $110 < X < 140$.

Given that a person is healthy (event H), a blood sugar measurement X is the Gaussian $(90, 20)$ random variable. Given that a person has diabetes, (event D), X is the Gaussian $(160, 40)$ random variable. A randomly chosen person is healthy with probability $P[H] = 0.9$ or has diabetes with probability $P[D] = 0.1$.

- ✓ (a) What is the conditional PDF $f_{X|H}(x)$?
- ✓ (b) Calculate the conditional probabilities $P[T^+|H]$, and $P[T^-|H]$.
- ✓ (c) Find $P[H|T^-]$, the conditional probability that a person is healthy given the event of a negative test.
- (d) When a person has an ambiguous test result (T^0), the test is repeated, possibly many times, until either a positive T^+ or negative T^- result is obtained. Let N denote the number of times the test is given. Assuming that for a given person the result of each test is independent of the result of all other tests, find the conditional PMF of N given event H that a person is healthy. Note that $N = 1$ if the person has a positive T^+ or negative T^- result on the first test.

$$\rightarrow f_{X|H}(x) = \mathcal{N}(90, 20)$$

$$f_{X|D}(x) = \mathcal{N}(160, 40)$$

$$P[H] = 0.9$$

$$P[D] = 0.1$$

$$a) f_{X|H}(x) = \frac{1}{\sqrt{2\pi} \cdot 20} e^{-\frac{(x-90)^2}{800}}, \quad -\infty < x < \infty$$

$$b) P[T^+|H] = \int_{140}^{\infty} \underbrace{f_{X|H}(x)}_{\mathcal{N}(90, 20)} dx$$

$$\begin{aligned} P(X > 140) &= P\left(\frac{X-90}{20} > \frac{140-90}{20}\right) \\ &= Q(2.5) \\ &= 0.0062 \end{aligned}$$

$$P(T^-|H) = \int_{-\infty}^{110} f_{X|H}(x) dx$$

$\hookrightarrow N(90, 20)$

$$= P(X < 110) = P\left(\frac{X-90}{20} < \frac{110-90}{20}\right)$$

$$= 1 - \Phi(1)$$

$$= 0.84$$

$$c) P(H|T^-) = \frac{P(T^-|H)P(H)}{P(T^-)}$$

$$= \frac{P(T^-|H)P(H)}{P(T^-|H)P(H) + P(T^-|D)P(D)}$$

$\begin{matrix} 0.84 & 0.9 \\ 0.84 & 0.1056 \end{matrix}$

$$P(T^-|H)P(H) + P(T^-|D)P(D)$$

$$P(T^-|D) = \int_{-\infty}^{110} f_{X|D}(x) dx$$

$\hookrightarrow N(160, 40)$

$$P(X \leq 110) = P\left(\frac{X-160}{40} \leq \frac{110-160}{40}\right)$$

$$= P(Z \leq -1.25)$$

$$= \Phi(1.25)$$

$$= 1 - \Phi(1.25)$$

$$= 1 - 0.8944$$

$$= 0.1056$$

$$= 0.84 \times 0.9$$

$$0.84 \times 0.9 + 0.1056 \times 0.1$$


$$= 0.756$$

$$= 756$$

$$0.756 + 0.011$$

$$= 767$$

$$= 0.9861$$



$$d) P(A) = P(110 \leq X \leq 140)$$

$$P(A|H) = P(110 \leq X \leq 140 | H)$$

↖ ambiguous

$$= P\left(\frac{20}{20} \leq \underbrace{\frac{X-90}{20}}_{Z \sim N(0,1)} \leq \frac{50}{20}\right)$$

↘ $N(90, 20)$

$$P(1 \leq Z \leq 2.5)$$

$$= Q(1) - Q(2.5)$$

$$= \Phi(2.5) - \Phi(1)$$

$$= 0.99379 - 0.84130$$

$$P(A|H) = 0.15249 = p$$

$$P_N(n) = p^{n-1} (1-p), \quad n = 1, 2, \dots$$

$$N \sim \text{Geometrik}(1-p)$$

$$E[N] = \frac{1}{1-p} \cong \frac{1}{0.85}$$

Problem 7.2.4

Y is the exponential (0.2) random variable. Given $A = \{Y < 2\}$, find:

(a) $f_{Y|A}(y)$,

(b) $E[Y|A]$.

$$f_Y(y) = 0.2 e^{-0.2y} \quad y > 0$$

$$a) f_{Y|A}(y) = \begin{cases} \frac{f_Y(y)}{P(A)} & y \in A \\ 0 & \text{o.w} \end{cases}$$

$$P(A) = P(Y < 2) = \int_0^2 0.2 e^{-0.2y} dy$$
$$= -e^{-0.2y} \Big|_0^2 = 1 - e^{-0.4}$$

$$f_{Y|A}(y) = \begin{cases} \frac{0.2 e^{-0.2y}}{1 - e^{-0.4}} & 0 \leq y \leq 2 \\ 0 & \text{o.w} \end{cases}$$

$$E[Y|A] = \int_{y \in A} y f_{Y|A}(y) dy$$

$$= \int_0^2 \frac{y \cdot 0.2 e^{-0.2y}}{1 - e^{-0.4}} dy$$

$$\left(\int u dv = uv - \int v du \right)$$

$$u = y \quad dv = 0.2 e^{-0.2y} dy$$

$$du = dy \quad v = -e^{-0.2y}$$

$$= \frac{1}{1-e^{-0.4}} \left[-y e^{-0.2y} \Big|_0^2 - \int_0^2 -e^{-0.2y} dy \right]$$

$$= \frac{1}{1-e^{-0.4}} \left[-2e^{-0.4} + \frac{-1}{0.2} e^{-0.2y} \Big|_0^2 \right]$$

$$= \frac{1}{1-e^{-0.4}} \left[-2e^{-0.4} - \frac{1}{0.2} e^{-0.4} + \frac{1}{0.2} \right]$$

Problem 7.2.6

The number of pages X in a document has PMF

$$P_X(x) = \begin{cases} 0.15 & x = 1, 2, 3, 4 \\ 0.1 & x = 5, 6, 7, 8 \\ 0 & \text{otherwise.} \end{cases}$$

A firm sends all documents with an even number of pages to printer A and all documents with an odd number of pages to printer B.

- Find the conditional PMF of the length X of a document, given the document was sent to A. What are the conditional expected length and standard deviation?
- Find the conditional PMF of the length X of a document, given the document was sent to B and had no more than six pages. What are the conditional expected length and standard deviation?

ör: 1,3,5,7 sayfa $\rightarrow$ B, 2,4,6,8 sayfa $\rightarrow$ A

$$P_{X|A}(x) = \begin{cases} \frac{P_X(x)}{P(A)} & , x \in A \\ 0 & \text{o.w} \end{cases}$$

$$a) P(A) = 0.15 + 0.15 + 0.1 + 0.1 = 0.5$$

$$P_{X|A}(x) = \begin{cases} \frac{0.15}{0.5} \rightarrow 0.3 & x = 2, 4, \\ \frac{0.1}{0.5} \rightarrow 0.2 & x = 6, 8 \\ 0 & \text{o.w} \end{cases}$$

$$b) \{B \cap (X \leq 6)\} = C = \{1, 3, 5\}$$

$$P(C) = P_X(1) + P_X(3) + P_X(5) = 0.4$$

$$P_{X|C}(x) = \begin{cases} \frac{0.15}{0.4} \rightarrow 3/8 & x = 1, 3 \\ \frac{0.1}{0.4} \rightarrow 2/8 & x = 5 \\ 0 & \text{o.w} \end{cases}$$

$$E[X|C] = 1 \times 3/8 + 3 \times 3/8 + 5 \times 2/8 = \frac{22}{8} = \frac{11}{4}$$

$$\text{Var}(X|C) = E[X^2|C] - (E[X|C])^2$$


$$E[X^2|C] = 1^2 \times 3/8 + 3^2 \times 3/8 + 5^2 \times 2/8 = \frac{80}{8} = 10$$

$$\text{Var}(X|C) = 10 - \left(\frac{11}{4}\right)^2 = 10 - \frac{121}{16} \approx 2.44$$

Problem 7.2.8

W is the Gaussian $(0, 4)$ random variable. Given the event $C = \{W > 0\}$, find the conditional PDF, $f_{W|C}(w)$, the conditional expected value, $E[W|C]$, and the conditional variance, $\text{Var}[W|C]$.

$$f_{W|C}(w) = \begin{cases} \frac{f_W(w)}{P(C)} & w \in C \\ 0 & \text{o.w.} \end{cases}$$



$$P(C) = 1/2$$

$$f_{W|C}(w) = \begin{cases} 2 \frac{1}{\sqrt{2\pi} 4} e^{-w^2/32} & w > 0 \\ 0 & \text{o.w.} \end{cases}$$

$$E[W|C] = \int_0^{\infty} 2w \frac{1}{\sqrt{2\pi} 4} e^{-w^2/32} dw$$

$$w^2 = u \quad 2w dw = du$$

$$w: 0 \rightarrow \infty$$

$$u: 0 \rightarrow \infty$$

$$= \int_{u=0}^{\infty} \frac{1}{4\sqrt{2\pi}} e^{-u/32} du = \left. \frac{-32}{4\sqrt{2\pi}} e^{-u/32} \right|_0^{\infty}$$

$$E[W|C] = \frac{8}{\sqrt{2\pi}}$$

$$E[W^2|C] = \int_{w=0}^{\infty} 2w^2 \frac{1}{4\sqrt{2\pi}} e^{-w^2/32} dw$$

$$= \int_{w=-\infty}^{\infty} w^2 \underbrace{\frac{1}{4\sqrt{2\pi}} e^{-w^2/32}}_{\mathcal{N}(0,4)} dw = 16$$

0 ortalamalı, 4 std sapmalı Gauss rast. değişkeni. varyansı (karesinin beklenen değeri)

$$\begin{aligned} \text{Var}(W|C) &= E[W^2|C] - E[W|C]^2 \\ &= 16 - \frac{64}{2\pi^2} = 16 - \frac{32}{\pi^2} = 12.75 \end{aligned}$$

Problem 7.2.9

The time between telephone calls at a telephone switch is the exponential random variable T with expected value $(0.01) \rightarrow \lambda = 10$

- (a) What is $E[T|T > 0.02]$, the conditional expected value of T ?
 (b) What is $\text{Var}[T|T > 0.02]$, the conditional variance of T ?

$$f_T(t) = 10e^{-10t}, \quad t \geq 0$$

$$a) A = \{T > 0.02\} \Rightarrow P(A) = \int_{0.02}^{\infty} 10e^{-10t} dt$$

$$f_{T|A}(t) = \begin{cases} \frac{10e^{-10t}}{e^{-0.2}}, & t > 0.02 \\ 0 & \text{o.w} \end{cases} = e^{-0.2}$$

$$E[T|A] = \int_{0.02}^{\infty} t \frac{10e^{-10t}}{e^{-0.2}} dt$$

$$\left(\int u dv = uv - \int v du \right)$$

$$\left\{ \begin{array}{ll} t = u & 10e^{-10t} dt = dv \\ dt = du & v = -e^{-10t} \end{array} \right.$$

$$= \left[-t e^{-10t} \Big|_{0.02}^{\infty} - \int_{0.02}^{\infty} -e^{-10t} dt \right] \frac{1}{e^{-0.2}}$$

$$= \left[0.02 e^{-0.2} + \frac{-1}{10} e^{-10t} \Big|_{0.02}^{\infty} \right] \frac{1}{e^{-0.2}}$$

$$= \left[0.02 e^{-0.2} + \frac{1}{10} e^{-0.2} \right] \frac{1}{e^{-0.2}} = \frac{e^{-0.2} (0.12)}{e^{-0.2}} = 0.12 //$$

Not: $0.12 = \underbrace{0.02}_{T > 0.02} + \underbrace{0.1}_{\frac{1}{\lambda}}$

HAFIZASIZ
MEMORYLESS
üstel dağılımın
özell.ğ.

b) $\text{Var}(T | T > 0.02)$

$$= E[T^2 | T > 0.02] + E[T | T > 0.02]^2$$

$$E[T^2 | T > 0.02] = \int_{0.02}^{\infty} t^2 \frac{10e^{-10t}}{e^{-0.2}} dt$$

2 defa kısmi integral.


Problem 7.2.10

As the final rider in the final 60 km time trial of the Tour de France, Roy must finish in time $T \leq 1$ hour to win the Tour. He has the choice of bike made of (1) carbon fiber or (2) titanium. On the carbon fiber bike, his speed V over the course is the continuous uniform random variable with $E[V] = 58$ km/hr and $\text{Var}[V] = 12$. On the titanium bike, V is the exponential random variable with $E[V] = 60$ km/hr. $f_{V|\text{Titan}}(v) = \frac{1}{60}e^{-v/60}$

- (a) Roy chooses his bike to maximize $P[W]$, the probability he wins the Tour. Which bike does Roy choose and what is $P[W]$?
- (b) Suppose instead that Roy flips a fair coin to choose his bike. What is $P[W]$?

$$P(T < 1 | \text{Carb}) \quad \begin{matrix} \text{Carb} \\ \text{Titan} \end{matrix} \quad P(T < 1 | \text{Titan})$$

$\frac{1}{3}$ $e^{-1} = \frac{1}{2.718}$



$$\frac{a+b}{2} = 58$$

$$\frac{(b-a)^2}{12} = 12$$

$$a+b = 116$$

$$b-a = 12$$

$$2b = 128$$

$$b = 64$$

$$a = 52$$

$$P(T < 1) = P\left(\frac{60}{V} < 1\right)$$

$$= P(V > 60)$$

$$P(V > 60 | \text{Carb}) = \frac{1}{3}$$

$$P(V > 60 | Titan) = \int_{60}^{\infty} \frac{1}{60} e^{-t/60} dt$$

$$= -e^{-t/60} \Big|_{60}^{\infty} = e^{-1}$$

b) Titanyum daha iyi ($\frac{1}{2.718}$ kazanma şansı)


$$P(V > 60) = \underbrace{P(V > 60 | Cerb)}_{\frac{1}{3}} \underbrace{P(Cerb)}_{\frac{1}{2}} + \underbrace{P(V > 60 | Titan)}_{\frac{1}{2.718}} \underbrace{P(Titan)}_{\frac{1}{2}}$$

$$P(V > 60) = \left(\frac{1}{3} + \frac{1}{2.718} \right) 0.5$$

Problem 7.3.1

X and Y are independent identical discrete uniform (1, 10) random variables. Let A denote the event that $\min(X, Y) > 5$. Find the conditional PMF $P_{X,Y|A}(x, y)$.

Sürekli ise




$$f_{X,Y}(x, y) = \underbrace{f_X(x)}_{1/9} \underbrace{f_Y(y)}_{1/9}$$

$$= \begin{cases} \frac{1}{81} & 1 < x, y < 10 \\ 0 & \text{o.w} \end{cases}$$

$$f_{X,Y|A}(x,y) = \begin{cases} \frac{\frac{1}{81} \xrightarrow{\frac{1}{25}}}{P(A)} & \begin{matrix} 1 < x,y < 10 \\ \min(x,y) > 5 \end{matrix} \\ 0 & \text{o.w} \end{cases}$$

$5 < x,y < 10$

$$P(A) = P(\min(X,Y) > 5) = P(X > 5) P(Y > 5) = \frac{5}{9} \times \frac{5}{9} = \frac{25}{81}$$



$$P_{X,Y}(x,y) = \begin{cases} \frac{1}{100} \\ 0 \end{cases} \quad x,y \in \{1, \dots, 10\}$$

o.w
 $x,y \in \{6, 7, 8, 9, 10\}$

$$P_{X,Y|A}(x,y) = \begin{cases} \frac{\frac{1}{100}}{P(A)} = \frac{1}{25} \\ 0 \end{cases}$$

o.w
 $x,y \in \{1, \dots, 10\}$
re
 $\min(x,y) \in \{6, 7, 8, 9, 10\}$
o.w

$$P(A) = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$$


$$P(A) = P(\min(X,Y) > 5) = P(X > 5) P(Y > 5)$$

$\frac{1}{2} \quad \frac{1}{2}$

Problem 7.3.5

X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} (x+y)/3 & 0 \leq x \leq 1; \\ & 0 \leq y \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$



Let $A = \{Y \leq 1\}$.

(a) What is $P[A]$?

(b) Find $f_{X,Y|A}(x,y)$.

(c) Find $f_{X|A}(x)$ and $f_{Y|A}(y)$.

a)

$$P[A] = \int_{y=0}^1 \int_{x=0}^1 f_{X,Y}(x,y) dx dy = \int_{y=0}^1 \int_{x=0}^1 \left(\frac{x}{3} + \frac{y}{3} \right) dx dy$$

$$= \int_{y=0}^1 \left(\frac{x^2}{6} + \frac{xy}{3} \Big|_0^1 \right) dy$$

$$= \int_0^1 \left(\frac{1}{6} + \frac{y}{3} \right) dy = \frac{y}{6} + \frac{y^2}{6} \Big|_0^1 = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$$

b)

$$f_{X,Y|A}(x,y) = \begin{cases} \frac{f_{X,Y}(x,y)}{P(A)} & 0 < x < 1 \\ & 0 < y < 1 \\ 0 & \text{o.w.} \end{cases}$$



$$f_{X,Y|A}(x,y) = \begin{cases} x+y & 0 < x < 1, \\ & 0 < y < 1, \\ 0 & \text{o.w.} \end{cases}$$

c)

$$f_{X|A}(x) = \int_{y=0}^1 f_{X,Y|A}(x,y) dy$$

$$f_{X|A}(x) = x + \frac{1}{2}, \quad 0 \leq x \leq 1$$

$$= \int_{y=0}^1 x+y dy = xy + \frac{y^2}{2} \Big|_0^1 = x + \frac{1}{2}$$



$$f_{Y|A}(y) = \int_{x=0}^1 f_{XY|A}(x, y) dx$$

$$= \int_{x=0}^1 (x+y) dx = \left. \frac{x^2}{2} + xy \right|_0^1$$

$$= \frac{1}{2} + y$$


$$\left. \begin{aligned} f_{XY|A} &= x+y \\ f_{X|A} &= \frac{1}{2} + x \\ f_{Y|A} &= \frac{1}{2} + y \end{aligned} \right\} X, Y \text{ bağımsız degil.}$$

Problem 7.3.9

X and Y are independent random variables with PDFs

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1, \\ 0 & \text{otherwise,} \end{cases}$$

$$f_Y(y) = \begin{cases} 3y^2 & 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$



Let $A = \{X > Y\}$.

- (a) What are $E[X]$ and $E[Y]$?
→ (b) What are $E[X|A]$ and $E[Y|A]$?

$$a) \quad E[X] = \int_0^1 \underbrace{2x}_{p^2 f} \cdot x dx = \left. \frac{2x^3}{3} \right|_0^1 = \frac{2}{3}$$

$$E[Y] = \int_0^1 3y^2 \cdot y dy = \left. \frac{3y^4}{4} \right|_0^1 = \frac{3}{4}$$

$$b) \quad f_{XY}(x, y) = 6xy^2 \quad 0 \leq x \leq 1, 0 \leq y \leq 1$$


$$f_{XY|A}(x, y) = \begin{cases} \frac{f_{XY|A}(x, y)}{P(A)} & 0 \leq y \leq x \leq 1 \\ 0 & \text{o.w.} \end{cases}$$

$P(A) \rightarrow 2/5$

$$f_{X,Y|A}(x,y) = \begin{cases} 15xy^2 & 0 \leq y \leq x \leq 1 \\ 0 & \text{o.w.} \end{cases}$$

$$f_{X|A}(x) = \int_0^x 15xy^2 dy = 5xy^3 \Big|_0^x = 5x^4, \quad 0 \leq x \leq 1$$

$$f_{Y|A}(y) = \int_y^1 15xy^2 dx = 15 \frac{x^2 y^2}{2} \Big|_y^1 = \frac{15y^2}{2} - \frac{15y^4}{2}, \quad 0 \leq y \leq 1$$



$$\int_0^1 5x^4 \cdot x dx = \frac{5x^6}{6} \Big|_0^1 = 5/6$$

$$E[Y|A] = \int_0^1 y \cdot \left(\frac{15y^2}{2} - \frac{15y^4}{2} \right) dy = \frac{15y^4}{8} - \frac{15y^6}{12} \Big|_0^1 = \frac{15}{8} - \frac{15}{12} = \frac{15}{24} = \frac{5}{8} //$$

$$\begin{aligned} E[Y|A] &= \int_{x=0}^1 \int_{y=0}^x y f_{X,Y|A}(x,y) dy dx \\ &= \int_{x=0}^1 \int_{y=0}^x 15xy^3 dy dx = \int_{x=0}^1 \left(\frac{15xy^4}{4} \Big|_0^x \right) dx \\ &= \int_0^1 \frac{15x^5}{4} dx = \frac{15x^6}{24} \Big|_0^1 \\ &= \frac{15}{24} = \frac{5}{8} // \end{aligned}$$

Problem 7.4.8

$Y = ZX$ where X is the Gaussian $(0, 1)$ random variable and Z , independent of X , has PMF

$$P_Z(z) = \begin{cases} 1-p & z = -1, \\ p & z = 1. \end{cases}$$

True or False:

- (a) Y and Z are independent. EVET
(b) Y and X are independent. HAYIR

$$f_{Y|Z}(y|z) \stackrel{?}{=} f_Y(y)$$

a) $f_{Y|Z=1}$ $Y=ZX$

$$P(Y \leq y | Z=1) = P(XZ \leq y | Z=1) \xrightarrow{\frac{d}{dx}} \\ = P(X \leq y | Z=1) = F_X(y) \rightarrow f_X(y) //$$

$$f_X(y) // \\ = f_{Y|Z=1}(y)$$

$$P(Y \leq y | Z=-1) = P(XZ \leq y | Z=-1)$$

$$= P(-X \leq y) = P(X \geq -y) \\ = 1 - P(X \leq -y) \\ = P(X > y)$$



$$\xrightarrow{\frac{d}{dy}} \{1 - F_X(-y)\}$$

$Z = -1$ de olsa $+1$ 'de olsa Y 'nin dağılımı, $f_{X|Z=-1} = f_X(y) //$ aynı olduğuna göre Y ve Z bağımsızdır.

b) Y ve X bağımsız mı?

$$f_Y(y) = \underbrace{f_{Y|Z=1}(y)}_{f_X(y)} \underbrace{P(Z=1)}_p + \underbrace{f_{Y|Z=-1}(y)}_{f_X(y)} \underbrace{P(Z=-1)}_{1-p}$$

$$f_Y(y) = f_X(y) \quad \text{V(0,1)}$$

$$\begin{aligned}
P(X \leq x, Y \leq y) &= P(X \leq x, XZ \leq y) \\
&= P(X \leq x, XZ \leq y | Z=1)P \\
&\quad + P(X \leq x, XZ \leq y | Z=-1)(1-p) \\
&= P(X \leq x, X \leq y)P + P(X \leq x, \overbrace{-X \leq y}^{x \geq -y})(1-p) \\
&= P(X \leq \min(x, y))P + P(-y \leq X \leq x)(1-p)P(x \geq -y)
\end{aligned}$$

Kısa Yol: X verildiğinde $Y=ZX$, ya $-x$ olur ya da $+x$ olur.

Normalde $N(0,1)$ dağılımı olan Y 'nin dağılımı, X verildiğinde değişti. Dolayısıyla bunlar bağımsız değildir

Problem 7.4.13


Packets arriving at an Internet router are either voice packets (v) or data packets (d). Each packet is a voice packet with probability p , independent of any other packet. Observe packets at the Internet router until you see two voice packets. Let M equal the number of packets up to and including the first voice packet. Let N equal the number of packets observed up to and including the second voice packet. Find the conditional PMFs $P_{M|N}(m|n)$ and $P_{N|M}(n|m)$. Interpret your results.

$$M \sim \text{Geometric}(p) \quad P_M(m) = (1-p)^{m-1} p, \quad m=1, \dots$$

$$P_N(n) = \binom{n-1}{1} \cdot (1-p)^{n-2} \cdot p \cdot p = \binom{n-1}{1} (1-p)^{n-2} p^2$$

$$P_{M|N}(m|n) = \frac{P(\text{ilk voice paketi } m\text{'inci de} \mid 2.\text{voice paketi } n\text{'inci de})}{n-1}, \quad m \leq n \quad \text{uniform}$$


Her zaman $N \geq M$



$$P_{M|N}(m|n) = \begin{cases} \frac{1}{n-1} & m=1, 2, \dots, n-1 \\ 0 & \text{o.w.} \end{cases}$$

UNIFORM

$$P_{N|M}(n|m) = P(n' \text{ incide 2. v.p.} \mid m' \text{ incide 1. v.p.})$$



$$P_{N|M}(n|m) = \begin{cases} (1-p)^{n-m-1} p & , n > m \\ 0 & \text{o.w.} \end{cases} \quad \left. \begin{array}{l} N-M \text{ Geometrik}(p) \\ \text{her v.p geldiğinde} \\ \text{süreç yenileniyor.} \end{array} \right\}$$

Problem 7.5.3

The probability model for random variable A is

$$P_A(a) = \begin{cases} 1/3 & a = -1, \\ 2/3 & a = 1, \\ 0 & \text{otherwise.} \end{cases}$$

The conditional probability model for random variable B given A is:

$$P_{B|A}(b|1) = \begin{cases} 1/3 & b = 0, \\ 2/3 & b = 1, \\ 0 & \text{otherwise,} \end{cases} \quad P_{B|A}(b|-1) = \begin{cases} 1/2 & b = 0, \\ 1/2 & b = 1, \\ 0 & \text{otherwise.} \end{cases}$$

- What is the probability model for random variables A and B ? Write the joint PMF $P_{A,B}(a,b)$ as a table.
- If $A = 1$, what is the conditional expected value $E[B|A = 1]$?
- If $B = 1$, what is the conditional PMF $P_{A|B}(a|1)$?
- If $B = 1$, what is the conditional variance $\text{Var}[A|B = 1]$ of A ?
- What is the covariance $\text{Cov}[A, B]$?

$$a) P_{B|A}(b|a) = \frac{P_{AB}(a,b)}{P_A(a)} \Rightarrow P_{AB}(-1,0) = \underbrace{P_{B|A}(0|-1)}_{1/3} \underbrace{P_A(-1)}_{1/3}$$

A \ B	0	1
-1	1/9	2/9
1	2/6	2/6

$$P_{AB}(-1,1) = \underbrace{P_{B|A}(1|-1)}_{2/3} \underbrace{P_A(-1)}_{1/3}$$

$$P_{AB}(1,0) = \underbrace{P_{B|A}(0|1)}_{1/2} \underbrace{P_A(1)}_{2/3}$$

$$P_{AB}(1,1) = \underbrace{P_{B|A}(1|1)}_{1/2} \underbrace{P_A(1)}_{2/3}$$

$$b) E[B|A=1]$$

$$= \sum_{b=0}^1 P_{B|A=1}(b) \cdot b$$

$$= 0 \cdot \underbrace{P_{B|A=1}(0)}_{1/2} + 1 \cdot \underbrace{P_{B|A=1}(1)}_{1/2}$$

$$= 1/2$$

$$c) E[A|B=1]$$

$$= \sum_{a=-1,1} a \cdot \underbrace{P_{A|B=1}(a)}$$

$$\frac{P(A=a, B=1)}{P(B=1)} = \frac{P(B=1|A=a) P(A=a)}{P(B=1)}$$

$$= \frac{-1 \cdot P(B=1|A=-1) P(A=-1)}{P(B=1)} + \frac{(+1) \cdot P(B=1|A=1) P(A=1)}{P(B=1)}$$

$$= \frac{(-1) \cdot \frac{2}{3} \cdot \frac{1}{3}}{5/9} + \frac{(+1) \cdot \frac{1}{2} \cdot \frac{2}{3}}{5/9}$$

$$= \frac{\frac{1}{3} - \frac{2}{9}}{5/9} = \frac{1}{5}$$

$$d) B=1 \text{ ise } \text{Var}(A|B=1) = E[A^2|B=1] - \underbrace{\left(E[A|B=1]\right)^2}_{1/5}$$

$$E[A^2|B=1] =$$

$$\frac{(-1)^2 \cdot \frac{2}{3} \cdot \frac{1}{3}}{5/9} + \frac{(+1)^2 \cdot \frac{1}{2} \cdot \frac{2}{3}}{5/9} = \frac{\frac{2}{9} + \frac{2}{6}}{5/9}$$

$$= 1 //$$

$$\text{Var}(A|B=1) = 1 - \left(\frac{1}{5}\right)^2 = 24/25$$

$$e) \text{Cov}(A, B) = E[AB] - E[A]E[B]$$

$$E[AB] = \cancel{-1 \cdot 0 \cdot \frac{1}{9}} + \cancel{(-1) \cdot (1) \cdot \frac{2}{9}} + \cancel{1 \cdot 0 \cdot \frac{2}{6}} + \cancel{1 \cdot 1 \cdot \frac{2}{6}} = \frac{2}{6} - \frac{2}{9} = \frac{1}{9}$$

$$E[A] = -1 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \frac{1}{3}$$

$$E[B] = 0 \cdot \frac{4}{9} + 1 \cdot \frac{5}{9} = \frac{5}{9}$$

$$\text{Cov}(A, B) = \frac{1}{9} - \frac{1}{3} \times \frac{5}{9} = -\frac{2}{27} //$$


977

Problem 7.5.6

Random variables X and Y have joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 1/2 & -1 \leq x \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) What is $f_Y(y)$?
(b) What is $f_{X|Y}(x|y)$?
(c) What is $E[X|Y=y]$?



$$a) f_Y(y) = \int_{-1}^y f_{X,Y}(x,y) dx = \int_{-1}^y \frac{1}{2} dx = \frac{y+1}{2}$$

$$b) f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \frac{\frac{1}{2}}{\frac{y+1}{2}} = \frac{1}{y+1}, \quad -1 \leq x \leq y$$

verilen y için x r.d. -1 ile y arasında düzgün dağılımlı. 0 o.w

$$c) E[X|Y=y] = \int_{x=-1}^y x f_{X|Y}(x) dx = \int_{-1}^y \frac{x}{y+1} dx$$

$$= \frac{x^2}{2(y+1)} \Big|_{-1}^y$$

$$= \frac{y^2 - 1}{2(y+1)} = \frac{y-1}{2} \quad -1 < y < 1$$

Problem 7.6.2

X and Y are jointly Gaussian random variables with $E[X] = E[Y] = 0$ and $\text{Var}[X] = \text{Var}[Y] = 1$. Furthermore, $E[Y|X] = X/2$. Find $f_{X,Y}(x,y)$.

$$X, Y \sim \mathcal{N}(0, 1)$$

$$f_{Y|X}(y|x) = \frac{1}{\tilde{\sigma}_Y \sqrt{2\pi}} e^{-\frac{(y - \tilde{\mu}_Y(x))^2}{2\tilde{\sigma}_Y^2}}$$

$$\tilde{\mu}_Y(x) = \mu_Y + \rho_{X,Y} \frac{\sigma_Y}{\sigma_X} (x - \mu_X)$$

$$\tilde{\sigma}_Y(x) = \sigma_Y \sqrt{1 - \rho_{X,Y}^2}$$

$$\rightarrow (\mu_Y = 0, \mu_X = 0, \sigma_Y = \sigma_X = 1)$$

$$\tilde{\mu}_Y(x) = \rho_{X,Y} x$$

$$E[Y|X] = \rho_{X,Y} X \rightarrow X/2$$

o zaman

$$\rho_{X,Y} = \frac{1}{2}$$

$$\begin{aligned} E[Y|X=1] &= 1/2 \\ E[Y|X=3] &= 3/2 \\ E[Y|X] &= X/2 \\ E[E[Y|X]] &= E[X] = 0 \end{aligned}$$

$$f_{X,Y}(x,y) = \frac{\exp\left[-\frac{x^2 - 2 \cdot \frac{1}{2}xy + y^2}{2\left(1 - \frac{1}{4}\right)}\right]}{2\pi \sqrt{1 - \frac{1}{4}}}$$

$$f_{X,Y}(x,y) = \frac{\exp\left[-\frac{x^2 - xy + y^2}{3/2}\right]}{2\pi \sqrt{3/4}}$$

Problem 10.1.1

$X_1, \dots, X_n$ is an iid sequence of exponential random variables, each with expected value 5.

- (a) What is $\text{Var}[M_9(X)]$, the variance of the sample mean based on nine trials?
- (b) What is $P[X_1 > 7]$, the probability that one outcome exceeds 7?
- (c) Use the central limit theorem to estimate $P[M_9(X) > 7]$, the probability that the sample mean of nine trials exceeds 7.

$$f_X(x) = \frac{1}{5} e^{-x/5}, \quad x \geq 0 \quad \boxed{\lambda = \frac{1}{5}}$$

$$\begin{aligned} \text{a) } \text{Var}(M_9(X)) &= \text{Var}\left(\frac{X_1 + X_2 + \dots + X_9}{9}\right) \\ &= \frac{1}{81} \cdot 9 \cdot \underbrace{\text{Var}(X)}_{\frac{1}{\lambda^2}} \\ &= \frac{1}{9} \times \frac{1}{25} \end{aligned}$$

$$\begin{aligned} \text{b) } P(X_1 > 7) &= \int_7^{\infty} \frac{1}{5} e^{-x/5} dx = -e^{-x/5} \Big|_7^{\infty} = e^{-7/5} \\ &\approx 0.25 \end{aligned}$$

c) Central Limit Theorem

$$\frac{X_1 + \dots + X_n}{n} \xrightarrow[n \rightarrow \infty]{\text{dist}} \mathcal{N}\left(E[X], \frac{\text{Var}(X)}{n}\right)$$

$$M_g(X) \sim \mathcal{N}\left(5, \frac{1}{9 \times 25}\right)$$

yaklaşık olarak ortalaması 5, ve

standart sapması $\frac{1}{15}$ olan bir Gauss


r.d. olarak modellenebilir.

$$\begin{aligned} P(M_g(X) > 7) &\approx P\left(\frac{M_g(X) - 5}{1/15} > \frac{7 - 5}{1/15}\right) \\ &\approx Q(30) \end{aligned}$$

Problem 10.1.2

$X_1, \dots, X_n$ are independent uniform random variables with expected value $\mu_X = 7$ and variance $\text{Var}[X] = 3$.

- What is the PDF of X_1 ?
- What is $\text{Var}[M_{16}(X)]$, the variance of the sample mean based on 16 trials?
- What is $P[X_1 > 9]$, the probability that one outcome exceeds 9?
- Would you expect $P[M_{16}(X) > 9]$ to be bigger or smaller than $P[X_1 > 9]$? To check your intuition, use the central limit theorem to estimate $P[M_{16}(X) > 9]$.



$$\mu_X = \frac{a+b}{2} = 7 \rightarrow a+b=14$$

$$\text{Var}(X) = \frac{(b-a)^2}{12} = 3$$

$$= b-a = 6$$

$$f_X(x) = \begin{cases} \frac{1}{6} & 4 \leq x \leq 10 \\ 0 & \text{o.w.} \end{cases}$$



$$\begin{cases} b=10 \\ a=4 \end{cases}$$

$$b) \text{Var}[M_{16}(X)] = \text{Var}\left[\frac{X_1 + \dots + X_{16}}{16}\right]$$

$$= \frac{\text{Var}(X)}{16} = \frac{3}{16}$$

$$c) P(X_1 > 9) = \frac{1}{6} (10 - 9) = \frac{1}{6}$$

d)



$$M_{16}(X) \sim \mathcal{N}\left(7, \frac{3}{16}\right)$$

$$P(M_{16}(X) > 9) = P\left(\frac{M_{16} - 7}{\sqrt{3/16}} > \frac{9 - 7}{\sqrt{3/16}}\right)$$

$$\approx Q\left(\frac{32}{3}\right) \text{ \textit{neredeyse inkenszt.}}$$

$$P(H < h) = P\left(\sqrt{\frac{X^2 + Y^2}{2}} < h\right)$$

$$= P(X^2 + Y^2 < 2h^2)$$

$$0 \leq r^2 < 2h^2 \quad = \iint \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}} dx dy$$

$$0 < \theta < 2\pi$$

$$\leftarrow (X^2 + Y^2 < 2h^2)$$

$$x = r \cos \theta \quad y = r \sin \theta \quad dx dy = r dr d\theta$$

$$x^2 + y^2 = r^2 \cos^2 \theta + r^2 \sin^2 \theta = r^2$$

Problem 10.2.1

The weight of a randomly chosen Maine black bear has expected value $E[W] = 500$ pounds and standard deviation $\sigma_W = 100$ pounds. Use the Chebyshev inequality to upper bound the probability that the weight of a randomly chosen bear is more than 200 pounds from the expected value of the weight.

$$P(|W - E[W]| > \varepsilon) \leq \frac{\text{Var}[W]}{\varepsilon^2}$$

Chebyshev Inequality

$$E[W] = 500$$

$$P(|W - E[W]| > 200) \leq \frac{100^2}{200^2} = \frac{1}{4}$$

Problem 10.2.2

For an arbitrary random variable X , use the Chebyshev inequality to show that the probability that X is more than k standard deviations from its expected value $E[X]$ satisfies

$$P[|X - E[X]| \geq k\sigma] \leq \frac{1}{k^2}.$$


For a Gaussian random variable Y , use the $\Phi(\cdot)$ function to calculate the probability that Y is more than k standard deviations from its expected value $E[Y]$. Compare the result to the upper bound based on the Chebyshev inequality.

Cheb. $P[|X - E[X]| \geq k\sigma] \leq \frac{\text{Var}[X]}{k^2 \sigma^2} = \frac{1}{k^2}$

$Y \sim N(\mu, \sigma)$

$$P[|Y - \mu| \geq k\sigma] = P(Y > \mu + k\sigma) + P(Y < \mu - k\sigma)$$

$$= P\left(\frac{Y - \mu}{\sigma} > \frac{k\sigma}{\sigma}\right) + P\left(\frac{Y - \mu}{\sigma} < -\frac{k\sigma}{\sigma}\right)$$



$$\begin{aligned}
 &= P(Z > k) + P(Z < -k) \\
 &= Q(k) + Q(k) \\
 &= 2Q(k)
 \end{aligned}$$

$k \setminus$	Cheb	Gauss
1	1	$2Q(1) = 2(1 - \Phi(1)) \approx 0.32$
2	$1/4$	$2Q(2) = 2(1 - \Phi(2)) \approx 0.054$
3	$1/9$	$2Q(3) = 2.7 \times 10^{-3}$
4	$1/16$	$2Q(4) = 6.34 \times 10^{-5}$

Problem 10.3.1

Let $X_1, X_2, \dots$ denote an iid sequence of random variables, each with expected value 75 and standard deviation 15.

- How many samples n do we need to guarantee that the sample mean $M_n(X)$ is between 74 and 76 with probability 0.99?
- If each X_i has a Gaussian distribution, how many samples n' would we need to guarantee $M_{n'}(X)$ is between 74 and 76 with probability 0.99?

$$E[M_n(X)] = 75$$

$$P[|M_n(X) - 75| > 1] \leq 0.01 \rightarrow \text{istere bu.}$$

$\downarrow$ $n=?$ $\downarrow$ $E[X]$

Chebyshev: $P[|M_n(X) - 75| > 1] \leq \frac{\text{Var}(X)}{n \cdot 1^2}$

$15^2 \leftarrow \frac{\text{Var}(X)}{n} \leq 0.01$

$$n > 100 \text{ Var}(X) = 15^2 \times 100 = 22500$$

Problem 10.3.2

Let X_A be the indicator random variable for event A with probability $P[A] = 0.8$. Let $\hat{P}_n(A)$ denote the relative frequency of event A in n independent trials.

(a) Find $E[X_A]$ and $\text{Var}[X_A]$.

(b) What is $\text{Var}[\hat{P}_n(A)]$?

(c) Use the Chebyshev inequality to find the confidence coefficient $1 - \alpha$ such that $\hat{P}_{100}(A)$ is within 0.1 of $P[A]$. In other words, find α such that

$$P[|\hat{P}_{100}(A) - P[A]| \leq 0.1] \geq 1 - \alpha.$$

(d) Use the Chebyshev inequality to find out how many samples n are necessary to have $\hat{P}_n(A)$ within 0.1 of $P[A]$ with confidence coefficient 0.95. In other words, find n such that

$$P[|\hat{P}_n(A) - P[A]| \leq 0.1] \geq 0.95.$$

$$X_A \sim \text{Bernoulli}(0.8)$$

$$\hat{P}_n(A)$$

$$a) E[X_A] = 1 \times 0.8 + 0 \times 0.2 = 0.8$$

$$\text{Var}[X_A] = p(1-p) = 0.8 \times 0.2 = 0.16$$

$$E[X_A^2] = 1^2 \times 0.8 + 0^2 \times 0.2 = 0.8$$

$$\text{Var}[X_A] = 0.8 - 0.8^2 = 0.16$$

$$\begin{aligned} b) \text{Var}[\hat{P}_n(A)] &= \text{Var}\left[\frac{X_{A_1} + X_{A_2} + \dots + X_{A_n}}{n}\right] \\ &= \frac{\text{Var}[X_A]}{n} = \frac{0.16}{n} \end{aligned}$$

$$c) \text{Chebyshev} \quad P[|X - E[X]| > \epsilon] \leq \frac{\text{Var}[X]}{n\epsilon^2}$$

$$\hat{P}_{100}(A) = \frac{\sum_{i=1}^{100} X_{A_i}}{n}$$

$$P[A] = E[X_A]$$

$$P\left[\left|\hat{P}_{100}(A) - P(A)\right| \leq 0.1\right] \geq 1 - \alpha$$

$$P\left[\left|\hat{P}_{100}(A) - P(A)\right| \geq 0.1\right] \leq \alpha$$

$$\text{Chebyshev} \rightarrow \frac{\text{Var}(X_A)}{100 \times 0.1^2}$$

$$\alpha = \frac{\text{Var}(X_A)}{100 \times 0.1^2} \rightarrow 0.16$$

$$\Rightarrow \boxed{\alpha = 0.16}$$

d)

$$\alpha = 0.05 = \frac{0.16}{n \times 0.1^2}$$

$$n = \frac{0.16}{0.05 \times 0.01} = 2500 \times 0.16 = 400$$

örnek
gerekir

Problem 10.4.1

$$X \sim \mathcal{N}(\mu, \sigma)$$

When X is Gaussian, verify Equation (10.24), which states that the sample mean is within one standard error of the expected value with probability 0.68. ✓

$$M_n(X) = \frac{X_1 + \dots + X_n}{n}$$

$$\text{Var}(M_n(X)) = \frac{\text{Var}(X)}{n} = \frac{\sigma^2}{n}$$

$M_n(X)$, $E[X]$ 'in tarafsız bir kestirisidir.

$$\left(E[M_n(X)] = E[X] \right)$$

Bu durumda $\text{MSE} = \text{Var}[M_n(X)] = \frac{\text{Var}(X)}{n} = \frac{\sigma^2}{n}$

$$\text{standard error} = \sqrt{E_n} = \frac{\sigma}{\sqrt{n}}$$

$$P\left(\left| \underset{\substack{\uparrow \\ \text{bestimm}}} M_n(x) - \underset{\substack{\downarrow \\ \text{gege}}} \mu \right| < \frac{\sigma}{\sqrt{n}}\right)$$

$$P\left(-\frac{\sigma}{\sqrt{n}} < M_n(x) - \mu < \frac{\sigma}{\sqrt{n}}\right)$$


$M_n(x) - \mu \Rightarrow$ beklenen dege sifiri.
varyans $\frac{\sigma^2}{n}$, std $\frac{\sigma}{\sqrt{n}}$

$$= P\left(-1 < \underbrace{\frac{M_n(x) - \mu}{\sigma/\sqrt{n}}}_{N(0,1)} < 1\right)$$

$$\Phi(1) - \Phi(-1) = \Phi(1) - (1 - \Phi(1))$$

$$= 2\Phi(1) - 1$$

$$= 2 \times 0.84 - 1 = \underline{\underline{0.68}}$$



$$\sigma^2 = 2\sigma^2 \rightarrow \sigma = 2$$

Problem 7.1.6


A random ECE student has height X in inches given by the PDF

$$f_X(x) = \frac{4e^{-(x-70)^2/8} + e^{-(x-65)^2/8}}{5\sqrt{8\pi} \cdot \sqrt{2\pi} \cdot 2}$$

(a) Sketch $f_X(x)$ over the interval $60 \leq x \leq 75$. (For purposes of sketching, note that $\sqrt{8\pi} \approx 5$)

(b) Find the probability that a random ECE student is less than 5 feet 8 inches tall.

(c) Use conditional PDFs to explain why $f_X(x)$ might be a reasonable model for ECE students.



$$b) P(X < 67) = P(X < 67 | K) P(K) + P(X < 67 | E) P(E)$$

$$K: \mu = 65, \sigma = 2 \Rightarrow P(X < 67 | K) = P\left(\frac{X - 65}{2} < \frac{67 - 65}{2}\right)$$

$$E: \mu = 70, \sigma = 2 \Rightarrow P(X < 67 | E) = P\left(\frac{X - 70}{2} < \frac{67 - 70}{2}\right)$$

$$= 0.2 \phi(1) + 0.8 \phi(-1.5)$$

Problem 7.2.4

Y is the exponential (0.2) random variable. Given $A = \{Y < 2\}$, find:

(a) $f_{Y|A}(y)$,

(b) $E[Y|A]$.

$$f_Y(y) = 0.2 e^{-0.2x}, \quad x > 0$$

$$a) f_{Y|A}(y) = \begin{cases} \frac{f_Y(y)}{P(A)} & 0 \leq y \leq 2 \\ 0 & \text{o.w} \end{cases}$$

$$P(A) = P(Y < 2) = \int_0^2 0.2 e^{-0.2x} dx \\ = F_Y(2) = 1 - e^{-0.2 \times 2} = 1 - e^{-0.4}$$

$$f_{Y|A}(y) = \begin{cases} \frac{0.2 e^{-0.2x}}{1 - e^{-0.4}} & 0 \leq y \leq 2 \\ 0 & \text{o.w} \end{cases}$$

$$b) E[Y|A]$$

$$= \int_0^2 y f_{Y|A}(y) dy = \int_0^2 \frac{y \cdot 0.2 e^{-0.2y}}{1 - e^{-0.4}} dy$$

$$\left(\begin{array}{l} 0.2 e^{-0.2y} dy = dv \\ -e^{-0.2y} = v \end{array} \quad \begin{array}{l} y = u \\ dy = du \end{array} \right)$$

$$\frac{1}{1 - e^{-0.4}} \left[-y e^{-0.2y} \Big|_0^2 - \int_0^2 -e^{-0.2y} dy \right]$$


$$\frac{1}{1-e^{-0.4}} \left[-2e^{-0.4} + -5e^{-0.2y} \right] \Big|_0^2$$

$$\frac{1}{1-e^{-0.4}} \left[-2e^{-0.4} + 5 - 5e^{-0.4} \right] = \frac{5 - 7e^{-0.4}}{1-e^{-0.4}}$$

Problem 7.2.8

W is the Gaussian $(0, 4)$ random variable. Given the event $C = \{W > 0\}$, find the conditional PDF, $f_{W|C}(w)$, the conditional expected value, $E[W|C]$, and the conditional variance, $\text{Var}[W|C]$.

$$P(C) = P(W > 0) = \frac{1}{2}$$



$$f_{W|C}(w) = \begin{cases} \frac{f_W(w)}{1/2} & w > 0 \\ 0 & w \leq 0 \end{cases}$$

$$E[W|C] = \int_0^{\infty} 2 \frac{1}{\sqrt{32\pi}} e^{-w^2/32} dw$$

$$\frac{w^2}{32} = u$$

$$\frac{w dw}{16} = du$$

$$E[W|C] = \int_0^{\infty} 2 \frac{1}{\sqrt{32\pi}} e^{-u} \frac{16}{1} du = \frac{32}{\sqrt{32\pi}} \int_0^{\infty} e^{-u} du$$

$$= \frac{32}{\sqrt{32\pi}}$$

$$\begin{aligned}
 E[W^2|C] &= \int_{-\infty}^{\infty} 2w^2 \frac{1}{\sqrt{2\pi}} e^{-w^2/32} dw \\
 &= \int_{-\infty}^{\infty} w^2 \underbrace{\frac{1}{\sqrt{2\pi}} e^{-w^2/32}}_{N(0,4)} dw = 16
 \end{aligned}$$

$$\begin{aligned}
 \text{Var}[W|C] &= E[W^2|C] - E[W|C]^2 \\
 &= 16 - \left(\frac{32}{\sqrt{32\pi}} \right)^2 \\
 &= 16 - \frac{32}{\pi}
 \end{aligned}$$

Problem 7.3.4

N and K have joint PMF

$$P_{N,K}(n,k) = \begin{cases} \frac{(1-p)^{n-1}p}{n} & n=1,2,\dots \\ & k=1,\dots,n, \\ 0 & \text{otherwise.} \end{cases}$$

Let B denote the event that $N \geq 10$.

(a) Find the conditional PMFs $P_{N|B}(n)$ and $P_{N,K|B}(n,k)$. Which should you find first?

(b) Find the conditional expected values $E[N|B]$, $E[K|B]$, $E[N+K|B]$, $\text{Var}[N|B]$, $\text{Var}[K|B]$, $E[NK|B]$.
 $E[N|B] + E[K|B]$

$$\begin{aligned}
 P(B) = P(N \geq 10) &= \sum_{n=10}^{\infty} \sum_{k=1}^n \frac{(1-p)^{n-1}p}{n} \\
 &= \sum_{n=10}^{\infty} \frac{(1-p)^{n-1}p}{n} \\
 &= (1-p)^9 p \left[1 + (1-p) + (1-p)^2 + \dots \right]
 \end{aligned}$$

$$a) \quad = (1-p)^9 p \quad \frac{1}{1-(1-p)} = (1-p)^9$$

$$P_N(n) = \sum_{k=1}^n \frac{(1-p)^{n-1} p}{n}$$

$$P_N(n) = (1-p)^{n-1} p, \quad n=1, \dots \Rightarrow \text{Geometric}(p)$$

$$P_{N|B}(n) = \begin{cases} \frac{P_N(n)}{P_B} & , \quad n=10, 11, \dots \\ 0 & n < 10 \end{cases}$$

$$= \begin{cases} \frac{(1-p)^{n-1} p}{(1-p)^9} & n=10, 11, \dots \rightarrow (1-p)^{n-10} p \\ 0 & \text{o.w.} \end{cases}$$

$$P_{N,K|B}(n, k) = \begin{cases} \frac{(1-p)^{n-1} p}{n (1-p)^9} & n=10, 11, \dots \\ 0 & \text{o.w.} \end{cases} \quad k=1, 2, \dots, n$$

$$b) E[N|B] = \sum_{n=10}^{\infty} n (1-p)^{n-10} p$$

$$n-10 = m-1 \Rightarrow m = n-9$$

$$= \sum_{m=1}^{\infty} (m+9) (1-p)^{m-1} p$$

$$= \underbrace{\sum_{m=1}^{\infty} m (1-p)^{m-1} p}_{\frac{1}{p}} + \underbrace{9 \sum_{m=1}^{\infty} (1-p)^{m-1} p}_{9 \cdot 1}$$

$$E[N|B] = \frac{1}{p} + 9$$

$$E[K|B] = ?$$

$$P_K(k) = \sum_{n=1}^{\infty} \frac{(1-p)^{n-1} p}{n}$$

$$E[K|B] = \sum_{k=1}^n k \sum_{n=10}^{\infty} P_{K,N|B}(n,k)$$

$$= \sum_{k=1}^n k \sum_{n=10}^{\infty} \frac{(1-p)^{n-1} p}{n (1-p)^9}$$

Problem 7.4.9

At the One Top Pizza Shop, mushrooms are the only topping. Curiously, a pizza sold before noon has mushrooms with probability $p = 1/3$ while a pizza sold after noon never has mushrooms. Also, a pizza is equally likely to be sold before noon as after noon. On a day in which 100 pizzas are sold, let N equal the number of pizzas sold before noon and let M equal the number of mushroom pizzas sold during the day. What is the joint PMF $P_{M,N}(m,n)$? Are M and N independent? Hint: Find the conditional PMF of M given N .

$N > M$ oldugh bellu

$$P_{M|N}(m|n) =$$

~~$$\frac{P(M=m, N=n)}{P(N=n)} = \frac{\binom{100}{n} \left(\frac{1}{2}\right)^n \left(\frac{1}{2}\right)^{100-n}}{\binom{100}{n} \left(\frac{1}{2}\right)^n \left(\frac{1}{2}\right)^{100-n}}$$~~

$$P_{M|N}(m|n) = \binom{n}{m} \left(\frac{1}{3}\right)^m \left(\frac{2}{3}\right)^{n-m}$$

$$\begin{aligned} P_{M,N}(m,n) &= P_{M|N}(m|n) P_N(n) \\ &= \binom{n}{m} \left(\frac{1}{3}\right)^m \left(\frac{2}{3}\right)^{n-m} \binom{100}{n} \left(\frac{1}{2}\right)^{100} \end{aligned}$$

Problem 7.5.5

Random variables N and K have the joint PMF

$$P_{N,K}(n,k) = \begin{cases} \frac{100^n e^{-100}}{(n+1)!} & n=0,1,\dots; \\ & k=0,1,\dots,n, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find the marginal PMF $P_N(n)$ and the conditional PMF $P_{K|N}(k|n)$.

(b) Find the conditional expected value $E[K|N=n]$.

(c) Express the random variable $E[K|N]$ as a function of N and use the iterated expectation to find $E[K]$.

a)

$$P_N(n) = \sum_{k=0}^n \frac{100^n e^{-100}}{(n+1)!} = \frac{100^n e^{-100}}{n!} \quad n=0,1,\dots$$

Poisson(100)

$$P_{K|N}(k|n) = \frac{P_{N,K}(n,k)}{P_N(n)} = \frac{\frac{100^n e^{-100}}{(n+1)!}}{\frac{100^n e^{-100}}{n!}} = \frac{1}{n+1}$$

$k=0,1,\dots,n$
Düzgün Dağılımı

b)

$$E[K|N=n] = \sum_{k=0}^n \frac{k}{n+1} = \frac{n}{2}$$

c)

$$E[K|N] = \frac{N}{2}$$

$$E[K] = \underbrace{E[E[K|N]]}_{\text{iterated expectation}} = E\left[\frac{N}{2}\right] = \frac{1}{2} E[N] = \frac{100}{2} = \underline{\underline{50}}$$